


**PART : PHYSICS**

1. A solid sphere and a hollow sphere are roll down purely equal distances on same inclined plane (starting from rest) in time  $t_1$  and  $t_2$  then

- (1)  $t_1 > t_2$                       (2)  $t_1 < t_2$                       (3)  $t_1 = 2t_2$                       (4)  $t_1 = t_2$

**Ans. (2)**

**Sol.**



$$a = \frac{g \sin \theta}{1 + \frac{I_{cm}}{mr^2}}$$

$$(I_{cm})_{solid} < (I_{cm})_{hollow}$$

$$a_{solid} > a_{hollow}$$

$$t_1 < t_2$$

2. A solid sphere rolls without slipping on a horizontal plane. What is ratio of translation kinetic energy to the rotation kinetic energy of the sphere?

- (1) 4/3                      (2) 3/4                      (3) 2/5                      (4) 5/2

**Ans. (4)**

**Sol.**  $V = R\omega$

$$\frac{K_t}{K_{rot}} = \frac{\frac{1}{2}mv^2}{\frac{1}{2} \times I\omega^2} = \frac{\frac{1}{2}mv^2}{\frac{1}{2} \times \frac{2}{5}mv^2} = \frac{5}{2}$$

3. Acceleration due to gravity on the surface of earth is  $g$  and acceleration due to gravity on a planet whose diameter is  $\frac{1}{3}$  of that of earth and same mass as that of earth is  $g'$ . If  $g' = ng$  then  $n$  is.

- (1) 9                      (2) 2                      (3)  $\frac{1}{2}$                       (4) 6

**Ans. (1)**

**Sol.**  $g = \frac{GM}{R^2}$

$$g' = \frac{GM}{\left(\frac{R}{3}\right)^2} = \frac{9GM}{R^2}$$

$$g' = 9g$$

4. If an object of rest mass  $M_0$  has momentum  $p$  and total energy  $E$  then which the of the following will be correct ? (where  $C$  is the velocity of light) -

(1)  $E^2 = M_0^2 C^2 + P^2 C^2$

(2)  $E^2 = M_0^2 C^4 + P^2 C^2$

(3)  $E = M_0 C^2 + PC^2$

(4)  $E^2 = M_0 C + PC$

Ans. (2)

Sol. In relativistic case  $m = \frac{m_0}{\sqrt{1 - \frac{v^2}{c^2}}}$

$$P^2 = \frac{m_0^2 v^2}{\left(1 - \frac{v^2}{c^2}\right)} = \frac{m_0^2 \frac{v^2}{c^2} c^2}{\left(1 - \frac{v^2}{c^2}\right)}$$

$$P^2 = \frac{m_0^2 c^2 \left(\frac{v^2}{c^2} - 1 + 1\right)}{\left(1 - \frac{v^2}{c^2}\right)}$$

$$P^2 = m_0^2 c^2 + \frac{m_0^2 c^2}{\left(1 - \frac{v^2}{c^2}\right)}$$

$$p^2 c^2 = m_0^2 c^2 + \left(\frac{m_0}{\sqrt{1 - \frac{v^2}{c^2}}}\right)^2 c^4$$


$$p^2 c^2 = -(m_0 c^2)^2 + (m c^2)^2$$

$$(m c^2)^2 = p^2 c^2 + (m_0 c^2)^2$$

$$E^2 = p^2 c^2 + m_0^2 c^4$$


$$E = \sqrt{(pc)^2 + (m_0 c^2)^2}$$

5. A bob of mass  $m$  is attached to a string of length ' $\ell$ '. If it is rotating in a horizontal circle of radius  $r$  with angular velocity  $\omega = \frac{3 \text{ rev}}{\pi \text{ sec}}$  and tension in the string is  $x(m\ell)$  then value of  $x$  is \_\_\_\_\_



Ans. 36.00

Sol.



$$\omega = \frac{3 \text{ rev}}{\pi \text{ sec}}$$

$$\omega = \frac{3}{\pi} \times 2\pi \frac{\text{rev}}{\text{sec}}$$





$$\omega = 6$$

$$T \sin \theta = m(\ell \sin \theta) \omega^2$$

$$T = m \omega^2 \ell = m(36) \ell = x(m \ell)$$

$$x = 36$$

6. Which of the following graph is correct. Hence F = Fahrenheit. & C = Celsius



Ans. (3)


Sol.  $\frac{C-0}{100} = \frac{F-32}{180} \Rightarrow C = \frac{5}{9}(F-32)$

$$C = \frac{5F}{9} - \frac{160}{9}$$

$$\text{Slope} = \frac{5}{9} = +\text{Ve}$$

$$\text{Intercept} = \frac{-160}{9}$$

7. An ideal gas is undergone through a cyclic process as shown in the graph. The net heat ejected by the gas during one cycle will be :-



- (1)  $5\pi\text{J}$                       (2)  $10\pi\text{J}$                       (3)  $15\pi\text{J}$                       (4)  $2.5\pi\text{J}$

**Ans. (1)**


**Sol.**  $W_{\text{cycle}} = \text{Area enclosed by P - V cycle} = \frac{\pi ab}{2}$

$$W_{\text{cycle}} = -\frac{1}{2} \pi (100 \times (10^{-2})^3) (100 \times 10^3)$$

$$Q_{\text{cycle}} = W_{\text{cycle}} = -5\pi$$

Heat rejected by the gas =  $5\pi\text{J}$

8.



Four charges each of value  $q_0$  are placed as shown. If potential energy of system is  $k_1$  in case (i) and PE of system is  $k_2$  in case (ii) then what is value of  $k_2 - k_1$

- (1)  $\frac{kq_0^2}{a} [3\sqrt{2} - 2]$                       (2)  $\frac{kq_0^2}{a} [5\sqrt{2} - 2]$                       (3)  $\frac{kq_0^2}{a} [3\sqrt{2} + 2]$                       (4) Zero

**Ans. (1)**

**Sol.**  $k_1 = 2 \left[ \frac{kq_0^2}{a} + \frac{kq_0^2}{a} + \frac{kq_0^2}{\sqrt{2}a} \right] = \frac{2kq_0^2}{a} \left[ 2 + \frac{1}{\sqrt{2}} \right]$

$$k_2 = 2 \left[ \frac{kq_0^2}{a/\sqrt{2}} \times 2 + \frac{kq_0^2}{a} \right] = \frac{2kq_0^2}{a} [2\sqrt{2} + 1]$$

$$k_2 - k_1 = \frac{kq_0^2}{a} (3\sqrt{2} - 2)$$

9. **Statement-1:** If in adiabatic process volume is decrease from V to V/2 then temperature also decreases  
**Statement-2:** Free expansion is irreversible as well as adiabatic  
 (1) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1.  
 (2) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1  
 (3) Statement-1 is True, Statement-2 is False  
 (4) Statement-1 is False, Statement-2 is True

**Ans. (4)**

**Sol.**  $T.V^{\gamma-1} = \text{constant}$   
 when volume decreases then temperature increases

10. Which of the following option is correct for increasing order of wave length

- (i) Infrared ray  
 (ii) x-ray  
 (iii) UV-ray  
 (iv) Microwave-ray

- (1) (i), (ii), (iii), (iv)      (2) (iv), (i), (iii), (ii)      (3) (ii), (iii), (i), (iv)      (4) (ii), (iii), (i), (iv)

**Ans. (3)**

**Sol.**  $\lambda_x < \lambda_{UV} < \lambda_{Ir} < \lambda_{Micro}$

11. Find the fringe width, if complete YDSE is immersed in a medium of refractive index  $\mu = 1.44$ .

Given  $\lambda_{air} = 690 \text{ nm}$ ,  $D = 0.72 \text{ meter}$        $d = 1.5 \text{ mm}$

- (1) 0.23 mm      (2) 1.23 mm      (3) 2.28 mm      (4) 0.40 mm

**Ans. (1)**

**Sol.** 
$$\beta_{red} = \frac{\lambda_{air}}{\mu} \frac{D}{d}$$

$$= \frac{690 \times 10^{-9} \times 72 \times 10^{-2}}{144 \times 10}$$

$$= \frac{690}{2 \times 3 \times 10^{-3}}$$

$$= 230 \times 10^{-6}$$
 0.23 mm

12. For which of the following inputs, the output will be zero (0) :-



- (A)  $x = 0, y = 0$   
 (C)  $x = 1, y = 0$

- (B)  $x = 0, y = 1$   
 (D)  $x = 1, y = 1$

(1) A,B,C


(2) B, C, D

(3) Only A

(4) Only D

**Ans. (2)**

Sol.



- (A)  $x = 0, y = 0 \Rightarrow \text{output} = 0 + 0 = 1$
- (B)  $x = 0, y = 1 \Rightarrow \text{output} = 0 + 1 = 0$
- (C)  $x = 1, y = 0 \Rightarrow \text{output} = 1 + 0 = 0$
- (D)  $x = 1, y = 1 \Rightarrow \text{output} = 1 + 1 = 0$

13. Power of two sources  $S_1$  and  $S_2$  are in ratio 2 : 1 and  $2 \times 10^{15}$  photons per sec of wavelength 600 nm from  $S_1$  are emitted then find the number of photons per second emitted from source  $S_2$  of wavelength 300 nm ?

- (1)  $5 \times 10^{15}$
- (2)  $2 \times 10^{15}$
- (3)  $5 \times 10^{14}$
- (4)  $2 \times 10^{14}$

Ans. (3)

Sol.  $P_1 = P = \frac{N_1 hc}{\lambda_1}$        $P = \frac{N_2 hc}{\lambda_2}$        $N \rightarrow \text{No. of photon/sec}$

$$P_2 = \frac{P}{2} = \frac{N_2 hc}{\lambda_2}$$

$$\frac{P_1}{P_2} = \frac{N_1}{\lambda_1} \cdot \frac{\lambda_2}{N_2}$$

$$N_2 = \frac{N_1 \lambda_2}{\lambda_1 \cdot 2} = \frac{2 \times 10^{15} \times 300}{600 \times 2}$$

$$n_2 = 5 \times 10^{14} \text{ per second}$$

14. **Statement (1)** : An electron in a uniform magnetic field, can move without changing its velocity vector.  
**Statement (2)** : In the above case, the magnetic field should be along the direction of its velocity.

- (1) Both statement 1 and statement 2 is correct, and statement 2 is the correct explanation of statement 1
- (2) Both statement 1 and Statement 2 is correct but statement 2 is not the correct explanation of statement 1
- (3) Statement 1 is correct, but statement 2 is incorrect
- (4) Statement 1 is incorrect, but statement 2 is correct.


Ans. (1)

15. A drone camera situated at a height of 18 km, capture an image of area  $400\text{km}^2$ , on a camera film of size  $2\text{cm} \times 2\text{cm}$ . find the focal length of the lens used in camera in mm.

- (1) 6
- (2) 14
- (3) 18
- (4) 27

Ans. (3)

Sol.



$$|m| = \frac{h_i}{h_o} = \frac{2\text{cm}}{20\text{km}} = \frac{2 \times 10^{-2}}{20 \times 10^3} = 10^{-6}$$

Since the image is real, so it will be inverted

$$m = -10^{-6} = \frac{-1}{10^6}$$

$$m = \frac{f}{f + u}$$

$$-\frac{10}{10^6} = \frac{f}{f + (-18\text{km})}$$

$$10^6 f = -f + 18 \text{ km}$$

$$10^6 f = 18 \text{ km}$$

$$f = \frac{18\text{km}}{10^6} = \frac{18 \times 10^3 \times 10^3 \text{mm}}{10^6}$$

$$f = 18 \text{ mm}$$

16. A spherical conductor carries a charge of  $4 \times 10^{-8}\text{C}$  brought in contact with an uncharged spherical conductor and they are separated by a distance  $r$ . Now force between them is  $9 \times 10^{-3}\text{ N}$ . Determine the separation between the charges.

(1) 7

(2) 4

(3) 9

(4) 2

Ans. (4)

Sol.




$$\frac{Kq^2}{R^2} = 9 \times 10^{-3}$$

$$\frac{9 \times 10^9 \times 2 \times 10^{-8} \times 2 \times 10^{-8}}{9 \times 10^{-3}} = R^2$$

$$R = 2 \text{ cm}$$

17.



In the following diagram polarizer  $P_1$  &  $P_2$  are orthogonal and  $P_3$  is aligned at  $45^\circ$  w.r.t.  $P_1$  and  $P_2$ . If unpolarised light of intensity  $I_0$  is incident on  $P_1$  and  $P_2$  and light after passing through  $P_3$  is used in YDSE.

at some point  $P$  where path difference is  $\frac{\lambda}{3}$ , What is resultant intensity?

- (1)  $\frac{I_0}{2}$                       (2)  $\frac{I_0}{3}$                       (3)  $\frac{I_0}{4}$                       (4)  $I_0$

Ans. (3)

Sol. Intensity after  $P_1$  &  $P_2$  is  $\frac{I_0}{2}$  and after  $P_3$   $\frac{I_0}{2} \cos^2 45 = \frac{I_0}{2} \left(\frac{1}{2}\right) = \frac{I_0}{4}$

$$\text{Now, } I_P = I_1 + I_2 + 2\sqrt{I_1 I_2} \cos \Delta\theta$$

$$= \frac{I_0}{4} + \frac{I_0}{4} + 2\frac{I_0}{4} \cos \left[ \frac{2\pi}{\lambda} \left( \frac{\lambda}{3} \right) \right]$$

$$= \frac{I_0}{2} + \frac{I_0}{2} \left[ -\frac{1}{2} \right]$$

$$I_P = \frac{I_0}{4}$$

18. A hot body is placed in the surrounding of temperature  $16^\circ\text{C}$ . During first 4 minutes, its temperature falls from  $40^\circ\text{C}$  to  $24^\circ\text{C}$ , then find its temperature after 4 minutes.

- (1)  $12^\circ\text{C}$                       (2)  $22^\circ\text{C}$                       (3)  $10^\circ\text{C}$                       (4)  $18.7^\circ\text{C}$

Ans. (4)

**Sol.**  $\left(\frac{dT}{dt}\right) = k(T - T_0)$





$$\left(\frac{40 - 24}{4 \text{ min}}\right) = k\left(\frac{24 + 40}{2} - 16\right) \quad \dots(i)$$

$$\left(\frac{24 - T}{4 \text{ min}}\right) = k\left(\frac{T + 24}{2} - 16\right) \quad \dots(ii)$$

Solving the equations we get

$$T = \frac{56}{3} = 18.7^\circ\text{C}$$

- 19.** An infinitely long wire has current 'i' and its radius is 'a'. Choose the correct graph for 'B' v/s 'r' where 'r' is distance from centre of wire



**Ans.** (2)

- 20.** The position vector of a particle varies with time as  $\vec{r} = (5t^2\hat{i} - 5t\hat{j})$  m. The magnitude and direction of velocity at  $t = 2$  will be ;

(1)  $5\sqrt{15}$

(2)  $5\sqrt{17}$  m/sec. at an angle of  $\tan^{-1}(4)$  with  $-y$  axis

(3)  $5\sqrt{17}$  m/sec. at an angle of  $\tan^{-1}(4)$  with  $x$  axis


(4)  $5\sqrt{17}$  m/sec. at an angle of  $\tan^{-1}(4)$  with  $-x$  axis

**Ans.** (2)

**Sol.**  $\vec{v} = \frac{d\vec{r}}{dt} = 10t\hat{i} - 5\hat{j}$

$$\vec{v}_{t=2} = 20\hat{i} - 5\hat{j} \Rightarrow |\vec{v}| = \sqrt{(20)^2 + (5)^2} = \sqrt{425}$$


$$= 5\sqrt{17} \text{ m/sec}$$



$$\tan\theta = \frac{20}{5} = 4$$

$$\theta = \tan^{-1}(4) \text{ with } -y \text{ axis}$$

- 21.** Find the difference of the current ( $I_A - I_B$ ). If  $n$  charge particles move in a circular path with  $\omega$  angular velocity and there are two ampere's loop are given



where  $I_A$  is net current passing through the amperian loop A and  $I_B$  is the net current passing through loop B.

(1)  $\frac{nq\omega}{2\pi}$

(2)  $\frac{nq\omega}{\pi}$

(3)  $\frac{2\pi\omega}{nq}$

(4)  $\frac{2\pi}{nq\omega}$

**Ans.** (1)

**Sol.** In loop B incoming and outgoing current is equal & opposite so  $I_B = 0$





$$T = \frac{2\pi}{\omega}$$

$$I_A = \frac{nq}{T}$$

$$I_A = \frac{nq\omega}{2\pi}$$

$$I_A - I_B = \frac{nq\omega}{2\pi}$$

22.  $x(t) = x_0 \sin^2\left(\frac{t}{2}\right)$  (where  $x_0 = 1$ ) find graph of Kinetic energy v/s  $x$



Ans. (3)

23. For the graph of stopping potential  $V_0$  v/s frequency which statement is correct .


- (1) graph is linear
- (2) slope is  $d/h$
- (3)  $h$  is related to slope
- (4) to find  $D$ , we don't require  $h$ .

Ans. (1)

Sol.  $eV_0 = h\nu - h\nu_0$

$$V_0 = \frac{h\nu}{e} - \frac{h\nu_0}{e}$$

$$V_0 = \frac{h\nu}{e} - \frac{d}{e}$$



$$\tan\theta = \frac{h}{e} = \text{slope}$$

24. A electron is moving in a circular path inside a solenoid with a time period of 75 ns. The current through the solenoid is 1 amp. Determine the number of turns per unit length of solenoid.



- (1)  $3.8 \times 10^3$
- (2)  $38 \times 10^3$
- (3)  $4.3 \times 10^3$
- (4)  $43 \times 10^3$

Ans. (1)

**Sol.** We know that

$$B = \frac{2\pi m}{qT} = \frac{2 \times 3.14 \times 9.1 \times 10^{-31}}{1.6 \times 10^{-16} \times 75 \times 10^{-9}}$$

$$B = 4.78 \times 10^{-3} \text{ T}$$

$$\text{So, } n = \frac{B}{\mu_0 I} = \frac{B}{\mu_0 I} \frac{4.78 \times 10^{-3}}{4\pi \times 10^{-2} \times 1}$$

$$n = 3.8 \times 10^3$$

**25.** A material has a bulk modulus of  $25 \times 10^{11} \text{ N/m}^2$ . If it undergoes a volumetric strain of 0.2%, what is the excess pressure applied ?

- (1)  $5 \times 10^8 \text{ N/m}^2$       (2)  $5 \times 10^9 \text{ N/m}^2$       (3)  $5 \times 10^{10} \text{ N/m}^2$       (4)  $5 \times 10^{11} \text{ N/m}^2$

**Ans.** (2)

**Sol.** The bulk modulus  $K$  of a material is defined by the formula :

$$K = \frac{\text{Excess Pressure}(P)}{\text{Volumetric Strain}(\Delta V/V)}$$

Given data :

$$\text{Bulk modulus, } K = 25 \times 10^{11} \text{ N/m}^2$$

$$\text{Volumetric strain, } \frac{\Delta V}{V} = 0.002$$

Rearranging the formula to solve for excess pressure :

$$P = K \times \left( \frac{\Delta V}{V} \right)$$

Thus, the excess pressure is :

$$5 \times 10^9 \text{ N/m}^2$$

## PART : CHEMISTRY

1. Arrange the following in ascending order wavelength

$\lambda_1$  = Infrared

$\lambda_2$  = Micro

$\lambda_3$  = X-ray

$\lambda_4$  = U.V.

(1)  $\lambda_3 < \lambda_4 < \lambda_1 < \lambda_2$       (2)  $\lambda_3 < \lambda_1 < \lambda_4 < \lambda_2$       (3)  $\lambda_2 < \lambda_1 < \lambda_4 < \lambda_3$       (4)  $\lambda_1 < \lambda_4 < \lambda_2 < \lambda_3$

**Ans. (1)**

**Sol.** Order of wavelength in EM spectrum:

Cosmic < Gamma < X-rays < UV < Visible < Intra Red < Micro < Radio

2.  $t_{2g}^3 e_g^1$  configuration is possible in:

(1) WFL; high spin      (2) WFL; low spin      (3) SFL; high spin      (4) SFL; low spin


**Ans. (1)**

**Sol.** WFL will not cause pairing & above is high spin arrangement (greater no. unpaired e<sup>-</sup>).

3. When ethane -1, 2-diammine is progressive added to aqueous of Nickel (II) chloride the sequence of colour changed observed will be

(1) Pale blue → Blue → Green → Violet      (2) Violet → Blue → Pale blue → Green  
 (3) Pale blue → Blue → Violet → Green      (4) Green → Pale blue → Blue → violet

**Ans. (4)**



**Sol.**

4. **Statement-I** :  $IE_1$  of Sn > Pb

**Statement-II** :  $IE_1$  of Si > Ge

(1) Both Statement I and statement II are true      (2) Both statement I and statement II are false  
 (3) Statement I is true but statement II is false      (4) Statement I is false but statement II is true

**Ans. (4)**

**Sol.** IE decreases down the gap. So  $SI > Ge$  & Pb, Sn exceptional

5. Compound  $\xrightarrow{\text{aquaregia}}$  B  $\xrightarrow[\text{CH}_3\text{COOH}]{\text{KNO}_2}$  Yellow ppt


(1) NiS      (2) ZnS      (3) CoS      (4) MnS

**Ans. (3)**



**Sol.**  $\text{CoS} \xrightarrow{\text{aquaregia}} \text{CoCl}_2 \xrightarrow[\text{CH}_3\text{COOH}]{\text{KNO}_2} \text{K}_3[\text{Co}(\text{NO}_2)_6] \downarrow \text{Yellow}$

6. 54.2% C, 9.2% H & 36.6% O are present in a compound. If its molar mass is 132 g, its molecular formula is  
 (1) C<sub>6</sub>H<sub>12</sub>O<sub>3</sub>                      (2) C<sub>4</sub>H<sub>8</sub>O<sub>2</sub>                      (3) C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>                      (4) None of these


Ans. (1)



7. Consider the following reactions





Calculate heat of reaction (kJ) for the given reaction



- (1)  $-(x + y)$                       (2)  $-(2x + y)$                       (3)  $x/y$                       (4)  $y - 2x$

Ans. (4)



$$\Delta H_r = \Delta H_1 + \Delta H_2$$

$$= -2x + y$$

$$= y - 2x \text{ kJ}$$

8. Match the following cations with respective spin magnetic moment

Ions	H(B.M)
(i) Ti <sup>3+</sup>	(p) 2.83
(ii) Sc <sup>+3</sup>	(q) 0.00
(iii) V <sup>+2</sup>	(r) 1.73
(iv) Ni <sup>+2</sup>	(s) 3.87

- (1) i-r; ii-q; iii-s; iv-p                      (2) i-p; ii-q; iii-r; iv-s  
 (3) i-s; ii-p; iii-q; iv-r                      (4) i-s; ii-p; iii-r; iv-q

Ans. (1)

9. Calculate the overall activation energy

$$K = \frac{k_1 k_3}{k_2} \quad E_{a_1} = 60 \text{ kJ}$$

$$E_{a_2} = 40 \text{ kJ}$$

$$E_{a_3} = 20 \text{ kJ}$$

Ans. (20)

**Sol.** 
$$e^{-E_a/RT} = \sqrt{\frac{e^{-E_{a1}/RT} \cdot e^{-E_{a3}/RT}}{e^{-E_{a2}/RT}}}$$

$$e^{-E_a/RT} = \sqrt{e^{\frac{(E_{a2} - E_{a1} - E_{a3})}{RT}}}$$

$$-E_a = (E_{a2} - E_{a1} - E_{a3}) \times \frac{1}{2}$$

$$E_a = (E_{a1} + E_{a3} - E_{a2}) \times \frac{1}{2}$$

$$E_a = \frac{1}{2}(60 + 20 - 40) = 20 \text{ kJ}$$

**10. Statement-I :** Oxygen-oxygen bond length in  $O_3$  is greater than  $O_2$ .

**Statement-II :** O-O bond order in  $O_3$  is 1.5 and O-O bond order in  $O_2$  is 2.

- (1) Both Statement I and statement II are true    (2) Both statement I and statement II are false  
 (3) Statement I is true but statement II is false    (4) Statement I is false but statement II is true

**Ans. (1)**

**11.** The successive ionisation energy (I.E.) of an element 'X' is given

	IE <sub>1</sub>	IE <sub>2</sub>	IE <sub>3</sub>	IE <sub>4</sub>	IE <sub>5</sub>
X →	500	600	2000	2200	2600

Data given in kJ/mol.


Find out the group number of element X.

- (1) Group-13                      (2) Group-14                      (3) Group-2                      (4) Group-13

**Ans. (3)**

**12.**  $MX_2$  observed molar mass: 65.6 Normal molar mass: 164

Find percentage dissociation.



**Ans. (78)**


**Sol.** 
$$i = \frac{\text{normal molar mass}}{\text{abnormal molar mass}} = \frac{164}{65.6} = 2.5$$





$$2.5 = 1 + 2\alpha$$

$$1.5 = 2\alpha$$

$$\alpha = 0.75$$


$$\% \alpha = 75$$

**13.**  Major Product?



Ans. (2)

Sol.



14. Match the following reactions given in Column-I with respective reagents given in Column-II.

Column-I		Column-II	
(a)	Etard Reaction	(p)	$\text{SnCl}_2 + \text{HCl}$
(b)	Gattermann Reaction	(q)	$\text{CrO}_2\text{Cl}_2$
(c)	Gattermann Koch Reaction	(r)	$\text{Cu} + \text{HCl}$
(d)	Staphen Reaction	(s)	$\text{CO} + \text{HCl}$ , Anhydrous $\text{AlCl}_3$

(1) a-(q); (b)-(r); (c)-(s); (d)-(p)


(2) a-(p); (b)-(q); (c)-(r); (d)-(s)

(3) a-(q); (b)-(s); (c)-(p); (d)-(r)

(4) a-(p); (b)-(r); (c)-(q); (d)-(s)

Ans. (1)

15.



Above conversion can be done by using which reagents among the following.

(1)  $\text{Fe}/\text{Br}_2$ ,  $\text{H}_2\text{O}(\Delta)$ ,  $\text{H}_2\text{SO}_4$


(2)  $\text{Ac}_2\text{O}$ ,  $\text{H}_2\text{SO}_4$ ,  $\text{Br}_2$ ,  $\text{NaOH}$

(3)  $\text{Ac}_2\text{O}$ ,  $\text{Br}_2/\text{AcOH}$ ,  $\text{H}_2\text{O}/\text{H}^+$

(4)  $\text{Ac}_2\text{O}$ ,  $\text{Br}_2/\text{Fe}$ ,  $\text{NaOH}$

Ans. (2)

16. Find the total number of  $sp$  and  $sp^2$  hybridised carbon atoms in the given compound.



Ans. (8)

Sol.  $sp^2 = 5$


$sp = 3$

Total = 8

17. In the carius method, 0.25 gm organic compound is heated with fuming  $\text{HNO}_3$  then  $\text{AgNO}_3$  is added it gives 0.15 gm  $\text{AgBr}$  if molecular mass of  $\text{AgBr}$  is 188 then find mass percentage of Br in that organic compound

Ans. (25.53%)

18.



Find total number of stereoisomers of the given compound.

Ans. (4)

19. Match the following Nitrogenous Bases with their respective structures.

**Column-I**


(a) Cytosine

(b) Uracil

(c) Guanine

(d) Adenine

**Column-II**



(1) a-(q); (b)-(r); (c)-(s); (d)-(p)

(2) a-(r); (b)-(s); (c)-(q); (d)-(p)

(3) a-(q); (b)-(s); (c)-(p); (d)-(r)

(4) a-(p); (b)-(r); (c)-(q); (d)-(s)

**Ans. (2)**

20. A hydrocarbon X that has molar mass 80 gm contains 90% carbon. Find degree of unsaturation in X.

**Ans. (3)**

## PART : MATHEMATICS

**1** If  $7 = 5 + \frac{1}{7} (5 + a) + \frac{1}{7^2} (5 + 2a) + \dots \infty$  term then the value of a is:

**Ans. (6)**

**Sol.**  $7 = 5 + \frac{1}{7} (5 + a) + \frac{1}{7^2} (5 + 2a) + \dots \infty$  ..... (i)

multiply with  $\frac{1}{7}$  ever term.

$$\frac{7}{7} = \frac{5}{7} + \frac{1}{7^2} (5 + a) + \dots \infty$$
 ..... (ii)

subtract from equation (i) .....equation (ii)

$$7 - \frac{7}{7} = 5 + \frac{1}{7} (a) + \frac{1}{7^2} (a) + \dots \infty$$

$$6 = 5 + \frac{1}{7} a [1 + \frac{1}{7} + \dots \infty]$$

$$1 = \frac{1}{7} a \left[ \frac{1}{1 - \frac{1}{7}} \right]$$

$$1 = \frac{1}{7} a \times \frac{7}{6}$$

$$a = 6$$

**2.** If A and B are binomial coefficients of 30<sup>th</sup> and 12<sup>th</sup> term of binomial expansion  $(1 + x)^{2n-1}$ . If  $2A = 5B$ , then the value of n is:

- (1) 19                                      (2) 20                                      (3) 21                                      (4) 40

**Ans. (3)**

**Sol.**  $T_{30} = {}^{2n-1}C_{29} \cdot x^{29}$ ,                       $A = {}^{2n-1}C_{29}$   
 $T_{12} = {}^{2n-1}C_{11} \cdot x^{11}$ ,                       $B = {}^{2n-1}C_{11}$

$$2A = 5B$$

$$\Rightarrow 2 \times {}^{2n-1}C_{29} = 5 \cdot {}^{2n-1}C_{11}$$

$$\Rightarrow \frac{2}{(2n-30)! \cdot 29!} = \frac{5}{(2n-12)! \cdot 11!}$$

$$n = 21.$$

**3.** In a arithmetic progression  $S_n$  represent the sum of n terms and  $S_{12} = 57$ ,  $S_{40} = 1030$  then the value of  $S_{30} - S_{10}$  is:

**Ans. (515)**

**Sol.**  $S_n = \frac{n}{2} [2a + (n-1)d]$

$$S_{12} = \frac{12}{2} [2a + 11d] = 57$$

$$2a + 11d = \frac{19}{2} \quad \dots(1)$$

$$S_{40} = \frac{40}{2} [2a + 39d] = 1030$$

$$2a + 39d = \frac{103}{2} \quad \dots(2)$$

Equation (2) - (1)

$$39d - 11d = \frac{103}{2} - \frac{19}{2}$$

$$d = \frac{3}{2}, a = \frac{-7}{2} \text{ now}$$

$$S_{30} - S_{10} = 15 [2a + 29d] - 5 [2a + 9d] = 20a + 390d$$

$$= 20 \times \left(\frac{-7}{2}\right) + 390 \times \left(\frac{3}{2}\right) = 515$$

4. Equation of the chord having mid point (3, 1) to the ellipse  $\frac{x^2}{25} + \frac{y^2}{16} = 1$  is :

(1)  $25x + 5y - 125 = 0$

(2)  $48x + 25y - 169 = 0$

(3)  $65x + 2y - 12 = 0$

(4)  $45x + 4y - 135 = 0$

Ans. (2)

Sol.  $S_1 = T$

$$= \frac{3^2}{25} + \frac{1^2}{16} - 1 = \frac{3x}{25} + \frac{y}{16} - 1$$

$$= \frac{9}{25} + \frac{1}{16} = \frac{3x}{25} + \frac{y}{16}$$

$$= 48x + 25y - 169 = 0$$

5. Let  $A = [a_{ij}]_{2 \times 2}$  such that  $a_{ij} \in \{0, 1\}$ . Probability that randomly chosen such matrix A is non-invertible is

(1)  $\frac{3}{8}$

(2)  $\frac{5}{8}$

(3)  $\frac{1}{2}$

(4)  $\frac{7}{8}$

Ans. (2)

Sol. Total number of matrices  $A = 2^4 = 16$

$$\text{Let } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

For non-invertible matrix  $|A| = 0$

$$\Rightarrow ad - bc = 0$$

$$ad = bc$$

Case 1  $\Rightarrow ad = bc = 0$

$$({}^2C_1 + {}^2C_2) ({}^2C_1 + {}^2C_2) = 9$$

Case 2  $\Rightarrow ad = bc = 1$

$$1 \times 1 = 1$$

$$\text{Required probability} = \frac{9+1}{16} = \frac{5}{8}$$

6. If system of equations  
 $x + 2y - 3z = 2$   
 $2x + \lambda y + 5z = 5$   
 $4x + 3y + \mu z = 33$   
 has infinite many solutions then  $\lambda + \mu$  is

- (1)  $\frac{244}{5}$                       (2)  $\frac{1334}{5}$                       (3)  $\frac{1296}{5}$                       (4)  $\frac{4997}{5}$

Ans. (2)

Sol.  $\Delta = 0 \quad \Delta = \begin{vmatrix} 1 & 2 & -3 \\ 2 & \lambda & 5 \\ 4 & 3 & \mu \end{vmatrix} = 0$

$12\lambda + \lambda\mu - 4\mu + 7 = \dots\dots\dots(i)$

$\Delta z = \begin{vmatrix} 1 & 2 & 2 \\ 2 & \lambda & 5 \\ 4 & 3 & 33 \end{vmatrix} = 0$


$\lambda = \frac{19}{5}$

from (i)  $\mu = 263$

$\lambda + \mu = \frac{19}{5} + 263 = \frac{1334}{5}$

7. The area bounded by  $y = e^x$ ,  $y = |e^x - 1|$  and y-axis is \_\_\_\_  
 (1)  $\ln 2$                       (2)  $1 + \ln 2$                       (3)  $1 - \ln 2$                       (4)  $2\ln 2$

Ans. (3)



$e^x = 1 - e^x$

$e^x = \frac{1}{2}$

$x = -\ln 2$

required area =  $\int_{-\ln 2}^0 (e^x - (1 - e^x)) dx = [2e^x - x]_{-\ln 2}^0$

$= 2 - (2e^{-\ln 2} + \ln 2) = 2 - 1 - \ln 2 = 1 - \ln 2$

8.  $f(x) = [x] + |x - 2|$  ;  $-2 < x < 3$   
 and  $m =$  number of points of discontinuity, and  
 $n =$  number of points of non differentiability.

Then the value of  $m + n$  is:

- (1) 8                      (2) 7                      (3) 9                      (4) - 10

Ans. (1)

**Sol.**  $f(x) = \begin{cases} -x & ; x \in (-2, -1) \\ 1-x & ; x \in [-1, 0) \\ 2-x & ; x \in [0, 1) \\ 3-x & ; x \in [1, 2) \\ x & ; x \in [2, 3) \end{cases}$

As we can see that function is discontinuous at  $-1, 0, 1$  and  $2$ .

Therefore, function is non-differentiable at  $-1, 0, 1$  and  $2$ .

Therefore,  $m = 4$  and  $n = 4$ .

Then,  $m + n = 8$ .

**9.** There is a group A of 5 boys and 3 girls and another group B of 5 boys and 6 girls. How many ways can we invite 4 boys and 4 girls for party with 5 from group A and 3 from group B.

- (1) 2850                      (2) 2550                      (3) 3150                      (4) 3450

**Ans. (3)**

**Sol.** Group A  $\Rightarrow$  5B, 3G

Group B  $\Rightarrow$  5B, 6G

4 Boys and 4 girls invite

5 from group A and 3 from group B

Case (i)	In group A	4B 1G	In B	0B 3G
	In group A	3B 2G	In B	1B 2G
	In group A	2B 3G	In B	2B 1G
	In group A	1B 4G	In B	3B 0G (not possible)

We have three cases than.

$${}^5C_4 \times {}^3C_1 \times {}^5C_0 \times {}^6C_3 = 5 \times 3 \times 1 \times 20 = 300$$

$${}^5C_3 \times {}^3C_2 \times {}^5C_1 \times {}^6C_2 = 10 \times 3 \times 5 \times 15 = 2250$$

$${}^5C_2 \times {}^3C_3 \times {}^5C_2 \times {}^6C_1 = 10 \times 1 \times 10 \times 6 = 600$$

$$300 + 2250 + 600 = 3150$$

**10.**  $2 \cos x \frac{dy}{dx} = \sin 2x - 2y \sin x$ ,  $y(x) = y$  and  $y(0) = 0$ , then find the value of  $y\left(\frac{\pi}{4}\right) + y\left(\frac{\pi}{4}\right)$  is:

- (1)  $\frac{1}{2}$                       (2)  $\frac{1}{\sqrt{2}}$                       (3)  $-\frac{1}{2}$                       (4)  $-\frac{1}{\sqrt{2}}$

**Ans. (2)**

**Sol.**  $(2 \cos x) y'(x) = \sin 2x - 2y(x) \sin x$

Put  $x = \frac{\pi}{4}$


$$2 \times \frac{1}{\sqrt{2}} y\left(\frac{\pi}{4}\right) = 1 - 2y\left(\frac{\pi}{4}\right) \times \frac{1}{\sqrt{2}}$$

$$y\left(\frac{\pi}{4}\right) + y\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}}$$

11. Number of real solutions of the equation  $x^2 + 3x + 2 = \min\{|x - 3|, |x + 2|\}$  is \_\_\_\_\_  
 (1) 0 (2) 1 (3) 2 (4) 3

Ans. (3)

Sol.



Total number of solutions = 2

12. Consider the differential equation  $x^2 \frac{dy}{dx} = 2xy + 3$  such that  $y(1) = 4$  then the value of  $2(y(2))$  is

Ans. (39)

Sol. Divide by  $x^2$

$$\frac{dy}{dx} - \frac{2}{x}y = \frac{3}{x^2}$$

$$\text{I.F.} = e^{\int -\frac{2}{x} dx} = e^{-2 \ln x} = e^{\ln x^{-2}} = \frac{1}{x^2}$$

$$y \cdot \text{I.F.} = \int \frac{3}{x^2} \cdot \text{I.F.} dx$$

$$\frac{y}{x^2} = 3 \int \frac{dx}{x^4}$$

$$\frac{y}{x^2} = \frac{-1}{x^3} + C$$

$$y(1) = 4 \Rightarrow 4 = -1 + C \Rightarrow C = 5$$

$$\frac{y}{x^2} = \frac{-1}{x^3} + 5$$

$$y = -\frac{1}{x} + 5x^2$$

$$y(2) = -\frac{1}{2} + 20 = \frac{39}{2}$$

$$2y(2) = 39$$

13. If

$$\lim_{x \rightarrow 0} \begin{vmatrix} a + \frac{\sin x}{x} & 1 & b \\ a & 1 + \frac{\sin x}{x} & b \\ a & 1 & b + \frac{\sin x}{x} \end{vmatrix} = \lambda a + \mu b + C$$

where  $\lambda$  and  $\mu$  are the coefficient of  $a$ ,  $b$  and  $c$  is constant then find the value of  $(\lambda + \mu + c)^2$

Ans. (16)

**Sol.**

$$\Rightarrow \begin{vmatrix} a+1 & 1 & b \\ a & 1+1 & b \\ a & 1 & b+1 \end{vmatrix} = \lambda a + \mu b + C$$

$$R_1 \rightarrow R_1 - R_2$$

$$R_2 \rightarrow R_2 - R_3$$

$$\begin{vmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ a & 1 & b+1 \end{vmatrix} = \lambda a + \mu b + C$$

$$C_2 \rightarrow C_1 + C_2$$

$$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ a & a+1 & b+1 \end{vmatrix} = \lambda a + \mu b + C$$

$$a + b + 2 = \lambda a + \mu b + C$$

$$\text{So } \lambda = 1, \mu = 1, C = 2$$

$$\text{Now, } (\lambda + \mu + C)^2 = 16$$

**14.** If  $f: (-\infty, \infty) \rightarrow (-\infty, 1)$ , and  $f(x) = \frac{2^x - 2^{-x}}{2^x + 2^{-x}}$ , then  $f(x)$  is

(1) one-one and onto

(2) one-one and into

(3) many-one and onto

(4) many-one and into

**Ans. (2)**

**Sol.**  $f(x) = \frac{2^{2x} - 1}{2^{2x} + 1} = \frac{2^{2x} + 1 - 2}{2^{2x} + 1} = 1 - \frac{2}{2^{2x} + 1}$

$2^{2x}$  is one-one so  $f(x)$  is one-one

$$\text{For } x \in (-\infty, \infty)$$

$$2^x \in (0, \infty)$$

$$2^{2x} + 1 \in (1, \infty)$$

$$\frac{1}{2^{2x} + 1} \in (0, 1)$$

$$\frac{-2}{2^{2x} + 1} \in (-2, 0)$$

$$1 - \frac{2}{2^{2x} + 1} \in (-1, 1)$$

Range of  $f(x)$  is  $(-1, 1)$  but codomain of  $f(x)$  is  $(-\infty, 1)$  so  $f(x)$  is into.

**15.** Given  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is a hyperbola with latus rectum  $12\sqrt{5}$  and eccentricity  $\sqrt{\frac{5}{2}}$  and another hyperbola

$\frac{x^2}{A^2} - \frac{y^2}{B^2} = -1$  with latus rectum  $15\sqrt{2}$ . If the product of transverse axis of both the hyperbola is  $100\sqrt{10}$ ,

eccentricity of the later hyperbola is:

(1) 0

(2) 2

(3)  $\sqrt{\frac{13}{5}}$

(4)  $\sqrt{\frac{11}{5}}$

**Ans. (4)**

**Sol.**  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

$$\frac{2b^2}{a} = 12\sqrt{5} \quad \dots\dots\dots (i)$$

$$\sqrt{1 + \frac{b^2}{a^2}} = \sqrt{\frac{5}{2}} \quad \dots\dots\dots (ii)$$

from (i) and (ii)

$$a = 4\sqrt{5} \text{ and } b^2 = 120$$

$$\frac{x^2}{A^2} - \frac{y^2}{B^2} = -1$$

$$\frac{2A^2}{B} = 15\sqrt{2} \quad \dots\dots\dots (iii)$$

(since product of transverse axis =  $100\sqrt{10}$ )

$$(2a).(2B) = 100\sqrt{10} \quad \dots\dots\dots (iv)$$

from (iii) & (iv)

$$A^2 = \frac{375}{4} \text{ and } B^2 = \frac{625}{8}$$

$$e_2 = \sqrt{1 + \frac{A^2}{B^2}} = \sqrt{\frac{11}{5}} \text{ (substituting the value of } A^2 \text{ and } B^2 \text{ from the above equation).}$$