

JEE (Main)-2026 Session-1
Question Paper with Solutions
(Mathematics, Physics, And Chemistry)
24 January 2026 Shift – 1

Time: 3 hrs.

M.M: 300

IMPORTANT INSTRUCTIONS:

- (1) The test is of 3 hours duration.
- (2) This test paper consists of 75 questions. Each subject (PCM) has 25 questions. The maximum marks are 300.
- (3) This question paper contains Three Parts. Part-A is Physics, Part-B is Chemistry and Part-C is Mathematics. Each part has only two sections: Section-A and Section-B.
- (4) Section - A: Attempt all questions.
- (5) Section - B: Attempt all questions.
- (6) Section - A (01 - 20) contains 20 multiple choice questions which have only one correct answer. Each question carries +4 marks for correct answer and -1 mark for wrong answer.
- (7) Section - B (21 - 25) contains 5 Numerical value-based questions. The answer to each question should be rounded off to the nearest integer. Each question carries +4 marks for correct answer and -1 mark for wrong answer.

Ans. (2)

Sol. For POI

$$2\lambda + 1 = 5\mu + 4; 3\lambda + 2 = 2\mu + 1; 4\lambda + 3 = \mu$$

$$\Rightarrow \lambda = \mu = -1$$

$$R(-1, -1, -1) \quad P(2\lambda + 1, 3\lambda + 2, 4\lambda + 3)$$

$$PR^2 = 29 \Rightarrow (2\lambda + 2)^2 + (3\lambda + 3)^2 + (4\lambda + 4)^2 = 29$$

$$\Rightarrow \lambda = 0 \text{ or } \lambda = -2 \text{ (Reject)}$$

$$\Rightarrow P(1, 2, 3)$$

$$Q(5\mu + 4, 2\mu + 1, \mu)$$

$$|PQ| = \sqrt{\frac{47}{3}} \Rightarrow PQ^2 = \frac{47}{3}$$

$$\Rightarrow (5\mu + 3)^2 + (2\mu - 1)^2 + (\mu - 3)^2 = \frac{47}{3}$$

$$\Rightarrow \mu = -\frac{1}{3}$$

$$Q = \left(\frac{7}{3}, \frac{1}{3}, -\frac{1}{3}\right)$$

$$(QR)^2 = \left(\frac{7}{3} + 1\right)^2 + \left(\frac{1}{3} + 1\right)^2 + \left(-\frac{1}{3} + 1\right)^2$$

$$= \frac{100 + 16 + 4}{9} = \frac{120}{9}$$

$$\Rightarrow 27 \times (QR)^2 = 27 \times \frac{120}{9} = 360$$

4. Let 729, 81, 9, 1, be a sequence and P_n denote the product of the first n terms of this sequence.

$$\text{If } 2 \sum_{n=1}^{40} (P_n)^{\frac{1}{n}} = \frac{3^\alpha - 1}{3^\beta} \text{ and } \gcd(\alpha, \beta) = 1, \text{ then}$$

$\alpha + \beta$ is equal to

$$(1) 73 \quad (2) 74$$

$$(3) 75 \quad (4) 76$$

Ans. (1)

Sol. $P_n = 729.81.9 \dots (n \text{ terms})$

$$= 3^6.3^4.3^2 \dots 3^{-2n+8}$$

$$P_n = 3^{6+4+2+\dots+(-2n+8)} = 3^{n(7-n)}$$

$$P_n^{1/n} = 3^{7-n}$$

$$\Rightarrow \sum_{n=1}^{40} (P_n)^{\frac{1}{n}} = 3^6 + 3^5 + \dots + (40 \text{ terms})$$

$$= 3^6 \left[\frac{1 - \left(\frac{1}{3}\right)^{40}}{1 - \frac{1}{3}} \right]$$

$$= \frac{3^6 [3^{40} - 1] \times 3^1}{3^{40} \times 2}$$

$$\sum (P_n)^{\frac{1}{n}} = \frac{(3^{40} - 1)}{2 \times 3^{33}}, \quad \alpha = 40$$

$$\beta = 33$$

$$\alpha + \beta = 73$$

5. Let $\vec{a} = 2\hat{i} + \hat{j} - 2\hat{k}$, $\vec{b} = \hat{i} + \hat{j}$ and $\vec{c} = \vec{a} \times \vec{b}$. Let \vec{d} be a vector such that $|\vec{d} - \vec{a}| = \sqrt{11}$, $|\vec{c} \times \vec{d}| = 3$ and the angle between \vec{c} and \vec{d} is $\frac{\pi}{4}$. Then $\vec{a} \cdot \vec{d}$

is equal to

$$(1) 11$$

$$(2) 3$$

$$(3) 0$$

$$(4) 1$$

Ans. (3)

$$\vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -2 \\ 1 & 1 & 0 \end{vmatrix}$$

$$\vec{c} = 2\hat{i} + 2\hat{k} + \hat{k}, |c| = 3$$

$$|\vec{c} \times \vec{d}| = 3$$

$$|\vec{c}| |\vec{d}| \sin \frac{\pi}{4} = 3 \Rightarrow |\vec{d}| = \sqrt{2}$$

$$|\vec{d} - \vec{a}| = \sqrt{11}$$

$$\Rightarrow |\vec{a}|^2 + |\vec{d}|^2 - 2\vec{a} \cdot \vec{d} = 11$$

$$9 + 2 - 2\vec{a} \cdot \vec{d} = 11$$

$$\vec{a} \cdot \vec{d} = 0$$

6. If the domain of the function

$$f(x) = \log_{(10x^2 - 17x + 7)}(18x^2 - 11x + 1)$$

is $(-\infty, a) \cup (b, c) \cup (d, \infty) - \{e\}$, then

$90(a + b + c + d + e)$ equals:

(1) 170

(2) 177

(3) 307

(4) 316

Ans. (4)

Sol. $18x^2 - 11x + 1 > 0$

$$(2x - 1)(9x - 1) > 0$$

$$x < \frac{1}{9} \text{ or } \frac{1}{2} < x$$

Also $10x^2 - 17x + 7 > 0$

$$(x - 1)(10x - 7) > 0$$

$$x < \frac{7}{10} \text{ or } 1 < x$$

& $10x^2 - 17x + 7 \neq 1$

$$x \in \left(-\infty, \frac{1}{9}\right) \cup \left(\frac{1}{2}, \frac{7}{10}\right) \cup (1, \infty) - \left\{\frac{6}{5}\right\}$$

$$90(a + b + c + d + e) = 90\left(\frac{1}{9} + \frac{1}{2} + \frac{7}{10} + 1 + \frac{6}{5}\right)$$

$$= 10 + 45 + 63 + 90 + 108 = 316$$

7. Let each of the two ellipses

$$E_1: \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, (a > b) \text{ and}$$

$$E_2: \frac{x^2}{A^2} + \frac{y^2}{B^2} = 1, (A < B) \text{ have eccentricity } \frac{4}{5}.$$

Let the lengths of the latus recta of E_1 and E_2 be ℓ_1 and ℓ_2 , respectively, such that $2\ell_1^2 = 9\ell_2$. If the distance between the foci of E_1 is 8, then the distance between the foci of E_2 is

(1) $\frac{96}{5}$

(2) $\frac{32}{5}$

(3) $\frac{16}{5}$

(4) $\frac{8}{5}$

Ans. (2)

Sol. $2ae = 8 \Rightarrow a = 5$

$$b^2 = a^2(1 - e^2)$$

$$b^2 = a^2 \times \frac{9}{25} \quad b^2 = 9$$

$$E_1: \frac{x^2}{25} + \frac{y^2}{9} = 1$$

$$\ell_1: \frac{2b^2}{a} = \frac{2 \times 9}{5} = \frac{18}{5}$$

$$A^2 = B^2(1 - e^2) \Rightarrow A^2 = \frac{9}{25}B^2 \Rightarrow A = \frac{3}{5}B$$

$$2\ell^2 = 9\ell_2 \Rightarrow 2\left(\frac{18}{5}\right)^2 = 9\ell_2 \Rightarrow \ell_2 = \frac{4 \times 18}{25}$$

$$\frac{2A^2}{B} = \frac{72}{25} \Rightarrow A^2 = \frac{36}{25}B$$

$$\frac{9}{25}B^2 = \frac{36B}{25} \Rightarrow B = 4,$$

$$\text{Distance between foci } 2Be = 2 \times \frac{4}{5} \times 4 = \frac{32}{5}$$

8. The value of $\frac{\sqrt{3} \operatorname{cosec} 20^\circ - \sec 20^\circ}{\cos 20^\circ \cos 40^\circ \cos 60^\circ \cos 80^\circ}$ is equal to

(1) 32

(2) 16

(3) 64

(4) 12

Ans. (3)

Sol.
$$E = \frac{\frac{\sqrt{3}}{\sin 20^\circ} - \frac{1}{\cos 20^\circ}}{\frac{1}{2} \cdot \frac{1}{4} \cdot \cos 60^\circ}$$

$$= \frac{(\sqrt{3} \cos 20^\circ - \sin 20^\circ)}{\cos 20^\circ \cdot \sin 20^\circ} \cdot 16$$

$$= \frac{\left(\frac{\sqrt{3}}{2} \cos 20^\circ - \frac{1}{2} \sin 20^\circ\right) 32 \times 2}{2 \cos 20^\circ \cdot \sin 20^\circ}$$

$$= \frac{\sin 40^\circ}{\sin 40^\circ} \times 64 = 64$$

9. Let $S = \left\{ z \in \mathbb{C} : \left| \frac{z-6i}{z-2i} \right| = 1 \text{ and } \left| \frac{z-8+2i}{z+2i} \right| = \frac{3}{5} \right\}$.

Then $\sum_{z \in S} |z|^2$ is equal to

- (1) 398 (2) 413
(3) 423 (4) 385

Ans. (4)

Sol. Solving $\left| \frac{z-6i}{z-2i} \right| = 1 \Rightarrow y = 4 \dots (1)$

(where $z = x + iy$)

Now solving $\left| \frac{z-8+2i}{z+2i} \right| = \frac{3}{5}$

$\Rightarrow x^2 + y^2 - 25x + 4y + 104 = 0 \dots (2)$

Solving (1) & (2) $\Rightarrow z = 17 + 4i$ & $8 + 4i$

$\Rightarrow \Sigma |z|^2 = (17)^2 + (4)^2 + (8)^2 + (4)^2 = 385$

10. If $\cot x = \frac{5}{12}$ for some $x \in \left(\pi, \frac{3\pi}{2} \right)$, then

$\sin 7x \left(\cos \frac{13x}{2} + \sin \frac{13x}{2} \right) +$

$\cos 7x \left(\cos \frac{13x}{2} - \sin \frac{13x}{2} \right)$ is equal to

- (1) $\frac{4}{\sqrt{26}}$ (2) $\frac{6}{\sqrt{26}}$
(3) $\frac{1}{\sqrt{13}}$ (4) $\frac{5}{\sqrt{13}}$

Ans. (3)

Sol. $\cot x = \frac{5}{12} \Rightarrow \cos x = \frac{-5}{13} = 2 \cos^2 \frac{x}{2} - 1$

$\cos \left(\frac{x}{2} \right) = -\frac{2}{\sqrt{13}}$ or $\frac{2}{\sqrt{13}}$ (rejected)

$\left\{ \because \frac{x}{2} \in \left(\frac{\pi}{2}, \frac{3\pi}{4} \right) \right\}$

$\left(\sin 7x \frac{\sin 13x}{2} + \cos 7x \frac{\cos 13x}{2} \right) + \left(\sin 7x \frac{\cos 13x}{2} - \cos 7x \frac{\sin 13x}{2} \right)$

$\cos \left(7x - \frac{13x}{2} \right) + \sin \left(7x - \frac{13x}{2} \right)$

$\cos \frac{x}{2} + \sin \left(\frac{x}{2} \right)$

$\frac{3}{\sqrt{13}} - \frac{2}{\sqrt{13}} = \frac{1}{\sqrt{13}}$

11. Let $f(t) = \int \left(\frac{1 - \sin(\log_e t)}{1 - \cos(\log_e t)} \right) dt, t > 1$.

If $f(e^{\pi/2}) = -e^{\pi/2}$ and $f(e^{\pi/4}) = \alpha e^{\pi/4}$, then α equals

- (1) $-1 - \sqrt{2}$
(2) $-1 - 2\sqrt{2}$
(3) $1 + \sqrt{2}$
(4) $-1 + \sqrt{2}$

Ans. (1)

Sol. $f(t) = \int \frac{1 - \sin(\ln t)}{1 - \cos(\ln t)} dt$

Let $\ln t = x \Rightarrow t = e^x \Rightarrow dt = e^x dx$

$= \frac{1}{2} \int \left(\operatorname{cosec}^2 \frac{x}{2} - 2 \cot \frac{x}{2} \right) e^x dx - t \cot \left(\frac{\ln t}{2} \right) + C$

$(\because \int (f(x) + f'(x))e^x dx = f(x) \cdot e^x + c)$

Now $f(e^{\pi/2}) = -e^{\pi/2} \cot \left(\frac{\pi}{4} \right) + C = -e^{\pi/2}$ (given)

$\Rightarrow C = 0$

Now $f(e^{\pi/4}) = -e^{\pi/4} \cot \left(\frac{\pi}{8} \right) + C = -e^{\pi/4}(\sqrt{2} + 1)$

12. Let R be a relation defined on the set $\{1, 2, 3, 4\} \times \{1, 2, 3, 4\}$ by

$R = \{((a, b), (c, d)) : 2a + 3b = 3c + 4d\}$.

Then the number of elements in R is

- (1) 6 (2) 18
(3) 12 (4) 15

Ans. (3)

Sol. (a, b)	(c, d)
(1, 1)	x
(1, 2)	x
(1, 3)	(1, 2)
(1, 4)	(2, 2)
(2, 1)	(1, 1)
(2, 2)	(2, 1)
(2, 3)	(3, 1)
(2, 4)	(4, 1)
(3, 1)	x
(3, 2)	x
(3, 3)	(1, 3)
(3, 4)	(2, 3)
(4, 1)	(1, 2)
(4, 2)	(2, 2)
(4, 3)	(3, 2)
(4, 4)	(4, 2)


13. Let A(1, 0), B(2, -1) and C($\frac{7}{3}, \frac{4}{3}$) be three points.

If the equation of the bisector of the angle ABC is $\alpha x + \beta y = 5$, then the value of $\alpha^2 + \beta^2$ is

- | | |
|--------|--------|
| (1) 8 | (2) 5 |
| (3) 13 | (4) 10 |

Ans. (4)

Sol.



$$\frac{BD}{DC} = \frac{AB}{AC} = \frac{\sqrt{2} \times 3}{5\sqrt{2}} = \frac{3}{5}$$

$$D = \left(\frac{12}{8}, \frac{4}{8} \right) = \left(\frac{3}{2}, \frac{1}{2} \right)$$

$$\text{Slope of AD} = \frac{-3/2}{\frac{1}{2}} = -3$$

$$3x + y = 5$$

$$\alpha = 3, \beta = 1; \alpha^2 + \beta^2 = 10$$

14. Let $S = \frac{1}{25!} + \frac{1}{3!23!} + \frac{1}{5!21!} + \dots$ up to 13 terms.

$$\text{If } 13S = \frac{2^k}{n!}, k \in \mathbb{N}, \text{ then}$$

$n + k$ is equal to

- | |
|--------|
| (1) 51 |
| (2) 52 |
| (3) 49 |
| (4) 50 |

Ans. (3)

$$\text{Sol. } \frac{1}{26!} \left(\frac{26!}{25!1!} + \frac{26!}{3!23!} + \frac{26!}{5!21!} + \dots + 13 \text{ terms} \right)$$

$$\frac{1}{26!} ({}^{26}C_1 + {}^{26}C_3 + {}^{26}C_5 + \dots + 13 \text{ terms})$$

$$\frac{1}{26!} ({}^{26}C_1 + {}^{26}C_5 + \dots + {}^{26}C_{25})$$

$$\Rightarrow S = \frac{1}{26!} \times 2^{25}$$

$$\Rightarrow 13S = \frac{2^{24}}{25!}$$

$$\text{so } n + k = 25 + 24 = 49$$

15. Consider an A.P.: a_1, a_2, \dots, a_n ; $a_1 > 0$. If $a_2 - a_1$

$$= \frac{-3}{4}, a_n = \frac{1}{4} a_1, \text{ and}$$

$$\sum_{i=1}^n a_i = \frac{525}{2}, \text{ then } \sum_{i=1}^{17} a_i \text{ is equal to}$$

- | | |
|---------|---------|
| (1) 476 | (2) 952 |
| (3) 238 | (4) 136 |

Ans. (3)

Sol. $S_n = \frac{n}{2}[a_1 + a_n] = \frac{525}{2}, d = \frac{-3}{4}$

$$\frac{n}{2}\left[a_1 + \frac{a_1}{4}\right] = \frac{525}{2}$$

$$\frac{5a_1n}{4} = 525$$

$$a_1n = 420$$

$$a_n = a_1 + (n-1)\left(\frac{-3}{4}\right)$$

$$\Rightarrow \frac{-3}{4}a_1 = \left(\frac{-3}{4}\right)(n-1) \Rightarrow a_1 = n-1$$

$$n(n-1) = 420$$

$$n^2 - n - 420 = 0$$

$$(n-21)(n+20) = 0$$

$$n = 21, a_1 = 20$$

$$\sum_{i=1}^{17} a_i = \frac{17}{2}[2a_1 + 16d]$$

$$= \frac{17}{2}\left[40 + 16\left(\frac{-3}{4}\right)\right]$$

$$= \frac{17}{2}[40 - 12]$$

$$= 17 \times 14 = 238$$

16. Let $\alpha, \beta \in \mathbb{R}$ be such that the function

$$f(x) = \begin{cases} 2\alpha(x^2 - 2) + 2\beta x & , x < 1 \\ (\alpha + 3)x + (\alpha - \beta) & , x \geq 1 \end{cases}$$

be differentiable at all $x \in \mathbb{R}$. Then $34(\alpha + \beta)$ is equal to

(1) 84 (2) 48

(3) 36 (4) 24

Ans. (2)

Sol. $f(x) = \begin{cases} 2\alpha x^2 + 2\beta x - 4\alpha & ; x < 1 \\ (\alpha + 3)x + \alpha - \beta & ; x \geq 1 \end{cases}$

$$f(1^+) = 2\alpha - \beta + 3, f(1^-) = -2\alpha + 2\beta$$

$$2\alpha - \beta + 3 = 2\beta - 2\alpha \Rightarrow 4\alpha - 3\beta + 3 = 0 \dots(1)$$

$$f'(1^+) = 4\alpha + 2\beta, f'(1^-) = \alpha + 3$$

$$4\alpha + 2\beta = \alpha + 3 \Rightarrow 3\alpha + 2\beta - 3 = 0 \dots(2)$$

Solving (1) & (2)

$$\text{We get } \alpha = \frac{3}{17}, \beta = \frac{21}{17}$$

$$\Rightarrow 34(\alpha + \beta) = 34 \times \frac{27}{17} = 48$$

17. From a lot containing 10 defective and 90 non-defective bulbs, 8 bulbs are selected one by one with replacement. Then the probability of getting at least 7 defective bulbs is :-

(1) $\frac{7}{10^7}$ (2) $\frac{81}{10^8}$

(3) $\frac{67}{10^8}$ (4) $\frac{73}{10^8}$

Ans. (4)

Sol. 10 defective & 90 non-defective

Req. probability = (7 def 1 fair) or (8 defective)

$$\text{Req. probability} = \frac{(10^7 \times 90) \times 8 + 10^8}{100^8}$$

$$= \frac{72 \times 10^8 + 10^8}{100^8} = \frac{73}{10^8}$$

18. The mean and variance of a data of 10 observations are 10 and 2, respectively. If an observations α in this data is replaced by β , then the mean and variance become 10.1 and 1.99, respectively. Then $\alpha + \beta$ equals.

(1) 10 (2) 15

(3) 5 (4) 20

Ans. (4)

Sol. Let first 10 numbers are $x_1, x_2, \dots, x_9, \alpha$

$$\Rightarrow \alpha + \sum_{i=1}^9 x_i = 100 \Rightarrow \sum_{i=1}^9 x_i = 100 - \alpha$$

$$\text{Variance} = \left(\frac{\sum x_i^2}{n} \right) - \left(\frac{\sum x_i}{n} \right)^2$$

$$\Rightarrow \frac{\sum x_i^2}{n} = 98$$

$$\Rightarrow x_1^2 + x_2^2 + \dots + x_9^2 + \alpha^2 = 1020 \Rightarrow \sum x_i^2 = 1020 - \alpha^2$$

In second case, let number are

$x_1, x_2, \dots, x_9, \beta$

$$100 - \alpha + \beta = 101 \quad \alpha - \beta + 1 = 0$$

$$\frac{\sum x_i^2 + \beta^2}{10} - (10.1)^2 = 1.99$$

$$\beta^2 - \alpha^2 = 20$$

$$\alpha = \frac{19}{2}$$

$$\beta = \frac{21}{2}$$


$$\alpha + \beta = \frac{19+21}{2} = 20$$

19. Let A_1 be the bounded area enclosed by the curves $y = x^2 + 2$, $x + y = 8$ and y -axis that lies in the first quadrant. Let A_2 be the bounded area enclosed by the curves $y = x^2 + 2$, $y^2 = x$, $x = 2$, and y -axis that lies in the first quadrant. Then $A_1 - A_2$ is equal to

(1) $\frac{2}{3}(2\sqrt{2} + 1)$ (2) $\frac{2}{3}(4\sqrt{2} + 1)$

(3) $\frac{2}{3}(\sqrt{2} + 1)$ (4) $\frac{2}{3}(3\sqrt{2} + 1)$

Ans. (1)




Sol.

$$A_1 = \int_0^2 ((8-x) - (x^2+2)) dx$$

$$= A_1 = \int_0^2 (6-x-x^2) dx$$

$$A_1 \left(6x - \frac{x^2}{2} - \frac{x^3}{3} \right)_0^2 = 12 - 2 - \frac{8}{3} = 10 - \frac{8}{3} = \frac{22}{3}$$



$$A_2 = \int_0^2 (x^2 + 2) dx - \frac{2}{3}(2\sqrt{2})$$

$$A_2 = \left(\frac{x^3}{3} + 2x \right)_0^2 - \frac{4\sqrt{2}}{3}$$

$$A_2 = \frac{8}{3} + 4 - \frac{4\sqrt{2}}{3} = \frac{20}{3} - \frac{4\sqrt{2}}{3}$$

$$A_1 - A_2 = \frac{2}{3} + \frac{4\sqrt{2}}{3}$$

20. The number of the real solutions of the equation :


$$x|x+3| + |x-1| - 2 = 0 \text{ is}$$

(1) 3 (2) 2

(3) 5 (4) 4

Ans. (1)

Sol.



(I)

$$x^2 + 3x + x - 1 - 2 = 0$$

$$x^2 + 4x - 3 = 0$$

$$x = -2 + \sqrt{7} \text{ (rejected)}, -2 - \sqrt{7} \text{ (rejected)}$$

(II)

$$x^2 + 3x + 1 - x - 2 = 0$$

$$x^2 + 2x - 1 = 0$$

$$x = -1 + \sqrt{2}, -1 - \sqrt{2}$$

(III)

$$-x^2 - 3x + 1 - x - 2 = 0$$

$$x^2 + 4x + 1 = 0$$

$$x = -2 - \sqrt{3}, -2 + \sqrt{3} \text{ (rejected)}$$

SECTION-B

21. Let a differentiable function f satisfy the equation

$$\int_0^{36} f\left(\frac{tx}{36}\right) dt = 4\alpha f(x).$$

If $y = f(x)$ is a standard parabola passing through the points $(2, 1)$ and $(-4, \beta)$,

Then β^α is equal to _____.

Ans. (64)

Sol. $\int_0^{36} f\left(\frac{tx}{36}\right) dt = 4\alpha f(x),$ Put $\frac{tx}{36} = y$

$$\frac{dy}{dt} = \frac{x}{36}$$

$$\int_0^x \frac{f(y)36dy}{x} = 4\alpha f(x)$$

$$\int_0^x f(y)dy = \frac{\alpha f(x)x}{9}$$

$$f(x) = \frac{\alpha}{9}(f(x) + xf'(x))$$

$$\left(1 - \frac{\alpha}{9}\right)f(x) = \frac{\alpha x}{9}f'(x) \Rightarrow (9 - \alpha)f(x) = \alpha x f'(x)$$

$$\frac{f'(x)}{f(x)} = \left(\frac{9}{\alpha} - 1\right)\frac{1}{x}$$

$$\log_e f(x) = \left(\frac{9}{\alpha} - 1\right)\log_e x + \log_e c$$

$$f(x) = cx^{\left(\frac{9}{\alpha} - 1\right)} \text{ for standard parabola}$$

$$\frac{9}{\alpha} - 1 = 2$$

$$\alpha = 3$$

$$f(x) = cx^2$$

passing through $(2, 1)$

$$1 = 4c \Rightarrow c = 1/4$$

$$y = \frac{x^2}{4} \text{ passing through } (-4, \beta)$$

$$\beta = 4$$

$$\beta^x = 4^3 = 64$$

22. Let a line L passing through the point $P(1, 1, 1)$ be perpendicular to the lines $\frac{x-4}{4} = \frac{y-1}{1} = \frac{z-1}{1}$

and $\frac{x-17}{1} = \frac{y-71}{1} = \frac{z}{0}$. Let the line L intersect

the yz -plane at the point Q . Another line parallel to L and passing through the point $S(1, 0, -1)$ intersects the yz -plane at the point R . Then the square of the area of the parallelogram $PQRS$ is equal to _____.

Ans. (6)

Sol. $d_1 = \langle 4, 1, 1 \rangle$ and $d_2 = \langle 1, 1, 0 \rangle$

$$d_L = d_1 \times d_2 = \begin{vmatrix} i & j & k \\ 4 & 1 & 1 \\ 1 & 1 & 0 \end{vmatrix} = \langle -1, 1, 3 \rangle$$

Line L passes through $P \langle 1, 1, 1 \rangle$ with $d_L = \langle -1, 1, 3 \rangle$

$$r(t) = \langle 1, 1, 1 \rangle + t \langle -1, 1, 3 \rangle$$

$$= \langle 1 - t, 1 + t, 1 + 3t \rangle$$

$$= \langle 1 - t, 1 + t, 1 + 3t \rangle$$

For point Q; $x = 0$

$$\Rightarrow t = 1$$

$$Q = \langle 0, 2, 4 \rangle$$

Another

Line parallel to L passes through $S \langle 1, 0, -1 \rangle$

with $d_L = \langle -1, 1, 3 \rangle$

$$r'(u) = \langle 1, 0, -1 \rangle + \mu \langle -1, 1, 3 \rangle$$

$$= \langle 1 - \mu, \mu, -1 + 3\mu \rangle$$

for point R, $x = 0$

$$\Rightarrow \mu = 1$$

$$R \langle 0, 1, 2 \rangle$$

Area of parallelogram with adjacent vectors \overrightarrow{PQ}

and \overrightarrow{PS}

$$\overrightarrow{PQ} = \langle -1, 1, 3 \rangle$$

$$\overrightarrow{PS} = \langle 0, -1, -2 \rangle$$

Area of parallelogram

$$\overrightarrow{PQ} \times \overrightarrow{PS} = \begin{vmatrix} i & j & k \\ -1 & 1 & 3 \\ 0 & -1 & -2 \end{vmatrix} = \langle 1, -2, 1 \rangle$$

$$\text{Area} = \sqrt{1^2 + (-2)^2 + 1^2} = \sqrt{6}$$

23. The number of numbers greater than 5000, less than 9000 and divisible by 3, that can be formed using the digits 0, 1, 2, 5, 9, if the repetition of the digits is allowed, is _____

Ans. (42)

Sol. (1) all different

$$5, 0, 1, 9 \Rightarrow \underline{3} = 6 \text{ ways}$$

(2) 2 alike, 2 different

$$5, 0, 0, 1 \Rightarrow 3 \text{ ways}$$

$$5, 1, 1, 2 \Rightarrow 3 \text{ ways}$$

$$5, 2, 2, 0 \Rightarrow 3 \text{ ways}$$

$$5, 2, 2, 9 \Rightarrow 3 \text{ ways}$$

$$5, 5, 0, 2 \Rightarrow 6 \text{ ways}$$

$$5, 5, 2, 9 \Rightarrow 6 \text{ ways}$$

$$5, 1, 9, 9 \Rightarrow 3 \text{ ways}$$

(3) 3 alike, 1 different

$$5, 5, 5, 0 \Rightarrow 3 \text{ ways}$$

$$5, 5, 5, 9 \Rightarrow 3 \text{ ways}$$

(4) 2 alike, 2 other alike

$$5, 5, 1, 1 \Rightarrow 3 \text{ ways}$$

Total ways = 42


24. Let $(2\alpha, \alpha)$ be the largest interval in which the function $f(t) = \frac{|t+1|}{t^2}$, $t < 0$, is strictly decreasing.

Then the local maximum value of the function

$g(x) = 2 \log_e(x-2) + \alpha x^2 + 4x - \alpha$, $x > 2$, is _____

Ans. (4)

Sol. Drawing graph of $f(t)$ for $t < 0$



$$g(x) = \log_e(x-2) - x^2 + 4x + 1; \quad x > 2$$

$$g'(x) = \frac{2}{x-2} - 2(x-2); \quad x > 2$$

$$g'(x) = \frac{1 - (x-2)^2}{(x-2)} = \frac{-(x-3)(x-1)}{(x-2)}$$

$$\begin{array}{c} + \quad - \\ \hline 2 \quad 3 \end{array} \quad \text{as } x > 2$$

maxima occur at $x = 3$

$$g(3) = 2 \log_e 1 - 9 + 12 + 1 = 4$$

25. The number of 3×2 matrices A , which can be formed using the elements of the set $\{-2, -1, 0, 1, 2\}$ such that the sum of all the diagonal elements of $A^T A$ is 5, is _____

Ans. (312)

Sol.
$$\begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \\ a_3 & b_3 \end{pmatrix}_{3 \times 2}$$

$$A^T A = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{pmatrix}_{2 \times 3} \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \\ a_3 & b_3 \end{pmatrix}_{3 \times 2}$$

$$= \begin{pmatrix} a_1^2 + a_2^2 + a_3^2 & \text{---} \\ \text{---} & b_1^2 + b_2^2 + b_3^2 \end{pmatrix}$$

$$\text{Tr}(A^T A) = a_1^2 + a_2^2 + a_3^2 + b_1^2 + b_2^2 + b_3^2 = 5$$

$$\{2, 1, 0, 0, 0, 0\}$$

$$\{2, -1, 0, 0, 0, 0\}$$

$$\{-2, 1, 0, 0, 0, 0\}$$

$$\{-2, -1, 0, 0, 0, 0\}$$

$$\{1, 1, 1, 1, 1, 0\}$$

$$\text{No. of ways} = \frac{6!}{4!} \times 4 + 2 \times \frac{6!}{5!} + 2 \times \frac{6!}{4!} + 2 \times \frac{6!}{3!2!}$$

$$= \frac{6!}{3!} + 2 \times 6 + 2 \times 15 \times 2 \times \frac{6!}{3!}$$

$$= 120 + 120 + 12 + 60 = 312$$

PHYSICS

SECTION-A

26. Match the List-I with List-II

List-I		List-II	
A.	Magnetic induction	I.	$MLT^{-2}A^{-2}$
B.	Magnetic flux	II.	$ML^2T^{-2}A^{-2}$
C.	Magnetic permeability	III.	$ML^0T^{-2}A^{-1}$
D.	Self inductance	IV.	$ML^2T^{-2}A^{-1}$

Choose the correct answer from the options given below:

- (1) A-IV, B-III, C-I, D-II (2) A-III, B-IV, C-II, D-I
 (3) A-I, B-III, C-IV, D-II (4) A-III, B-IV, C-I, D-II

Ans. (4)

Sol. Magnetic induction

$$F = qvB$$

$$[B] = \left[\frac{F}{qV} \right]$$

$$[B] = [MT^{-2}A^{-1}]$$

Magnetic Flux (ϕ)

$$\phi = (B) \cdot (\text{Area})$$

$$[\phi] = [ML^2T^{-2}A^{-1}]$$

Magnetic Permeability

$$[\mu] = [MLT^{-2}A^{-2}]$$

Self inductance

$$\text{Using } U = \frac{1}{2} LI^2$$

$$[\text{Self inductance}] = [ML^2T^{-2}A^{-1}]$$


A – III, B – IV, C – I, D – II

27. Three charges $+2q$, $+3q$ and $-4q$ are situated at $(0, -3a)$, $(2a, 0)$ and $(-2a, 0)$ respectively in the xy plane. The resultant dipole moment about origin is _____

- (1) $2qa(3\hat{j} - \hat{i})$ (2) $2qa(3\hat{i} - 7\hat{j})$
 (3) $2qa(7\hat{i} - 3\hat{j})$ (4) $2qa(3\hat{j} - 7\hat{i})$

Ans. (3)

Sol.



$$\vec{p} = q_1\vec{r}_1 + q_2\vec{r}_2 + q_3\vec{r}_3$$

$$\vec{p} = (2q)(-3a)\hat{j} + (3q)(2a)\hat{i} + (-4q)(-2a)\hat{i}$$


$$\vec{p} = 2qa(7\hat{i} - 3\hat{j})$$

28. A cylindrical block of mass M and area of cross section A is floating in a liquid of density ρ and with its axis vertical. When depressed a little and released the block starts oscillating. The period of oscillation is _____.

- (1) $2\pi\sqrt{\frac{M}{\rho Ag}}$
 (2) $\pi\sqrt{\frac{2M}{\rho Ag}}$
 (3) $\pi\sqrt{\frac{\rho A}{Mg}}$
 (4) $2\pi\sqrt{\frac{\rho A}{Mg}}$

Ans. (1)

Sol.



At equilibrium

$$\rho Ahg = Mg$$

After displacing by x ,

$$Ma = -\rho A(h+x)g + Mg$$

$$Ma = -\rho Ahg - \rho Axg + Mg$$

$$Ma = -\rho Axg$$

$$a = \left(\frac{-\rho Ag}{M} \right) x$$

on comparing with,

$$a = -\omega^2 x$$

$$\omega = \sqrt{\frac{\rho Ag}{M}}$$

$$T = \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{M}{\rho Ag}}$$

29. Density of water at 4°C and 20°C are 1000 kg/m^3 and respectively. The increase in internal energy of 4 kg water when it is heated from 4°C to 20°C is ___ J.

(Specific heat capacity of water = 4.2 J/kg . and 1 atmospheric pressure = 10^5 Pa)

- (1) 315826.2
 (2) 234699.2
 (3) 258700.8
 (4) 268799.2

Ans. (4)

Sol. $Q = mS\Delta T = 4 \times 4200 \times 16 \text{ J} = 268800 \text{ J}$

$$W = P\Delta V$$

$$\Delta V = \left(\frac{m}{\rho_f} - \frac{m}{\rho_i} \right) = 4 \left[\frac{1}{998} - \frac{1}{1000} \right]$$

$$P = 10^5 \text{ Pa.}$$


$$\therefore W = 10^5 \times 4 \times \left[\frac{1}{998} - \frac{1}{1000} \right] = \frac{8 \times 10^5}{10^3 \times 998} \approx 0.8 \text{ J}$$

$$\Delta U = Q - W = 268799.2 \text{ J}$$

30. Two resistors of 100Ω each are connected in series with a 9V battery. A voltmeter of 400Ω resistance is connected to measure the voltage drop across one of the resistors. The voltmeter reading is ___ V.

- (1) 3 (2) 4.5
 (3) 4 (4) 2

Ans. (3)



Sol.

Current in circuit.

$$I = \frac{E}{R_{eq}}$$

$$R_{eq} = 100 + \frac{400 \times 100}{400 + 100} = 180\Omega$$

$$\therefore I = \frac{9}{180} = \frac{1}{20} \text{ A}$$


$$\text{Reading of voltmeter} = V = I \times 80 = \frac{1}{20} \times 80 = 4\text{V}$$

31. The exit surface of a prism with refractive index n is coated with a material having refractive index $\frac{n}{2}$. When this prism is set for minimum angle of deviation it exactly meets the condition of critical angle. The prism angle is _____


- (1) 60° (2) 15°
 (3) 30° (4) 45°

Ans. (1)

Sol. $i = e$ & $r = A/2$ for minimum deviation



For TIR ; $r = \theta_c$



$$\sin r = \sin \theta_c$$

$$\sin r = \frac{n/2}{n}$$

$$\sin r = \frac{1}{2}$$

$$\sin \frac{A}{2} = \sin 30^\circ$$

$$\frac{A}{2} = 30^\circ$$

$$A = 60^\circ$$

32. Two electrons are moving in orbits of two hydrogen like atoms with speeds 3×10^5 m/s and 2.5×10^5 m/s respectively. If the radii of these orbits are nearly same then the possible order of energy states are _____ respectively.

- (1) 6 and 5 (2) 9 and 8
 (3) 8 and 10 (4) 10 and 12

Ans. (1)

Sol. $V \propto \frac{Z}{n}$
 $r \propto \frac{n^2}{Z}$

Thus ; $r \propto \frac{n}{V}$

Radii are same then

$$\frac{n_1}{V_1} = \frac{n_2}{V_2}$$

$$\frac{n_1}{n_2} = \frac{3 \times 10^5}{2.5 \times 10^5} = \frac{6}{5}$$

Possible order is 6 and 5

33. In a microscope of tube length 10 cm two convex lenses are arranged with focal length of 2 cm and 5 cm. Total magnification obtained with this system for normal adjustment is $(5)^k$. The value of k is _____

- (1) 2 (2) 5
 (3) 3.5 (4) 4


Ans. (1)

Sol. $f_o = 2$ cm, $f_e = 5$ cm

$$l = 10$$
 cm, $D = 25$ cm

$$M = \frac{l}{f_o} \cdot \frac{D}{f_e} = 25$$

34. For the series LCR circuit connected with 220 V, 50 Hz a.c source as shown in the figure, the power factor is $\frac{\alpha}{10}$. The value of α is _____



- (1) 4 (2) 10
 (3) 6 (4) 8

Ans. (3)

Sol. Power factor = $\frac{R}{Z}$

$$Z = \sqrt{R^2 + (X_L - X_C)^2}$$

$$= \sqrt{60^2 + (150 - 70)^2} = 100\Omega$$

$$\therefore \text{Power factor} = \frac{60}{100} = \frac{6}{10}$$

Then a = 6

35. Match the **List-I** with **List-II**

List-I		List-II	
A	Radio-wave	I	is produced by Magnetron valve
B	Micro-wave	II	Due to change in the vibrational modes of atoms
C	Infrared-wave	III	Due to inner shell electrons moving from higher energy level to lower energy level
D	X-ray	IV	Due to rapid acceleration of electrons

Choose the **correct** answer from the options given below:

- (1) A-II, B-IV, C-III, D-I (2) A-IV, B-III, C-I, D-II
 (3) A-IV, B-I, C-II, D-III (4) A-IV, B-II, C-I, D-III

Ans. (3)

Sol. Radio wave \Rightarrow Produced by rapid acceleration of electrons

Micro wave \Rightarrow By magnetron valve

Infrared wave \Rightarrow Change in vibrational modes

X ray \Rightarrow Transition of inner shell electrons from high energy level to low energy level.


A - IV, B - I, C - II, D - III

36. A spring of force constant 15 N/m is cut into two pieces. If the ratio of their length is 1:3, then the force constant of smaller piece is ___ N/m

- (1) 15 (2) 20
 (3) 60 (4) 45

Ans. (3)

Sol.




$$Kl = \text{constant}$$

$$Kl = K' \left(\frac{l}{4} \right)$$

$$K' = 4K$$

$$K' = 60 \text{ N/m}$$

37. Two resistors 2Ω and 3Ω are connected in the gaps of bridge as shown in figure. The null point is obtained with the contact of jockey at some point on wire XY. When an unknown resistor is connected in parallel with 3Ω resistor, the null point is shifted by 22.5 cm toward Y. The resistance of unknown resistor is ___ Ω .



- (1) 3 (2) 2
 (3) 4 (4) 1

Ans. (2)

Sol. Initially, $\frac{2}{3} = \frac{x}{100 - x}$

$$\Rightarrow x = 40 \text{ cm}$$

Now when 'R' connected in parallel

$$\frac{2}{3+R} = \frac{40 + 22.5}{60 - 22.5} = \frac{62.5}{37.5}$$

$$\therefore R = 2\Omega$$

$$0.02 = \frac{(400 - 350)(10^{-3})g}{(400 + 350)(10^{-3}) + \frac{I}{R^2}}$$

$$\frac{I}{R^2} = \frac{50 \times 10^{-3} g}{0.02} - 750 \times 10^{-3} = 23.75$$

$$I = 23.75 \times 4 \times 10^{-4} = 9.5 \times 10^{-3} \text{ kg-m}^2$$

42. An unpolarised light is incident at an interface of two dielectric media having refractive indices of 2 (incident medium) and $2\sqrt{3}$ (medium) respectively. To satisfy the condition that reflected and refracted rays are perpendicular to each other, the angle of incidence is _____.

- (1) 60° (2) 10° (3) 30° (4) 45°

Ans. (1)

Sol. Brewster's law

$$\tan\theta = \mu_{\text{rel}} = \sqrt{3}$$

$$\theta = 60^\circ$$


43. A boy thrown a ball into air at 45° from the horizontal to land it on a roof of a building of height H. If the ball attains maximum height in 2 s and lands on the building in 3 s after launch, then value of H is _____ m.

$$(g = 10 \text{ m/s}^2)$$

- (1) 20 (2) 10 (3) 25 (4) 15

Ans. (4)

Sol. $T = \frac{2u_y}{g} = 4$



$$\Rightarrow u_y = \frac{40}{2} = 20 \text{ m/s}$$

$$y = u_y \Delta t - \frac{1}{2} g (\Delta t)^2$$

$$\Rightarrow H = 20 \times 3 - 5 \times 9$$

$$= 60 - 45$$

$$= 15 \text{ m}$$

44. There are three co-centric conducting spherical shells A, B and C of radii a, b and c respectively. The potential of the spheres A, B and C respectively, are :


(1) $\frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{a} \right), \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{b} \right), \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{c} \right)$

(2) $\frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{a} \right), \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{b} + \frac{q_3}{c} \right), \frac{1}{4\pi\epsilon_0} \left(\frac{q_1}{a} + \frac{q_2}{b} + \frac{q_3}{c} \right)$

(3) $\frac{1}{4\pi\epsilon_0} \left(\frac{q_1}{a} + \frac{q_2}{b} + \frac{q_3}{c} \right), \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{b} + \frac{q_3}{c} \right), \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{c} \right)$

(4) $\frac{1}{4\pi\epsilon_0} \left(\frac{q_1}{a} + \frac{q_2}{b} + \frac{q_3}{c} \right), \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{b} \right), \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{c} \right)$

Ans. (3)



Sol.

$$V_A = \frac{Kq_1}{a} + \frac{Kq_2}{b} + \frac{Kq_3}{c} = \frac{1}{4\pi\epsilon_0} \left(\frac{q_1}{a} + \frac{q_2}{b} + \frac{q_3}{c} \right)$$

$$V_B = \frac{Kq_1}{b} + \frac{Kq_2}{b} + \frac{Kq_3}{c} = \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2}{b} + \frac{q_3}{c} \right)$$

$$V_C = \frac{Kq_1}{c} + \frac{Kq_2}{c} + \frac{Kq_3}{c} = \frac{1}{4\pi\epsilon_0} \left(\frac{q_1 + q_2 + q_3}{c} \right)$$

45. Three masses 200 kg, 300 kg and 400 kg are placed at the vertices of an equilateral triangle with sides 20 m. They are rearranged on the vertices of a bigger triangle of side 25 m and with the same centre. The work done in this process _____ J.

(Gravitational constant $G = 6.7 \times 10^{-11} \text{ N m}^2 / \text{kg}^2$)

(1) 9.86×10^{-6}


(2) 2.85×10^{-7}

(3) 1.74×10^{-7}

(4) 4.77×10^{-7}

Ans. (3)

Sol



Work done by external agent :

$$W_{\text{ext}} = \Delta U$$

$$U_i = -\frac{Gm_1m_2}{r_i} - \frac{Gm_2m_3}{r_i} - \frac{Gm_1m_3}{r_i} : r_i = 20 \text{ m}$$

$$U_f = -\frac{Gm_1m_2}{r_f} - \frac{Gm_2m_3}{r_f} - \frac{Gm_1m_3}{r_f} : r_f = 25 \text{ m}$$

$$U_i = \frac{-6.67 \times 10^{-11}}{20} [200 \times 300 + 300 \times 400 + 200 \times 400]$$

$$= \frac{-6.67 \times 10^{-11}}{20} \times 26 \times 10^4 = -86.71 \times 10^{-8} \text{ J}$$

$$U_f = \frac{-6.67 \times 10^{-11}}{0.25} [200 \times 300 + 300 \times 400 + 200 \times 400]$$

$$= \frac{-6.67 \times 10^{-11}}{0.25} \times 26 \times 10^4 = -693.68 \times 10^{-9}$$

$$= -69.36 \times 10^{-8} \text{ J}$$

$$\Delta U = U_f - U_i = 1.74 \times 10^{-7} \text{ J}$$

SECTION-B

46. A short bar magnet placed with its axis at 30° with an external field of 800 Gauss, experiences a torque of 0.016 N.m. The work done in moving it from most stable to most unstable position is $\alpha \times 10^{-3}$ J. The value of α is _____.

Ans. (64)

$$\text{Sol. } \tau = \mu B \sin \theta \Rightarrow 0.016 = \mu \times B \times \frac{1}{2}$$

$$\Rightarrow \mu = \frac{0.032}{B}$$

$$W_{\text{ext}} = U_f - U_i = \mu B - (\mu B) = 2\mu B$$

$$= 2 \times \frac{0.032}{B} \times B$$

$$= 0.064 \text{ J}$$

47. A gas of certain mass filled in a closed cylinder at a pressure of 3.23 kPa has temperature 50°C . The gas is now heated to double its temperature. The modified pressure is _____ Pa.

Ans. (3730)

Sol. As per NTA

$$V = \text{constant}$$

$$\text{so } P \propto T$$

$$T_i = 50^\circ\text{C} = 323 \text{ K}$$

$$T_f = 100^\circ\text{C} = 373 \text{ K}$$

$$\Rightarrow \frac{P_f}{P_i} = \frac{T_f}{T_i}$$

$$\Rightarrow \frac{P_f}{3.23 \text{ kPa}} = \frac{373}{323}$$

$$\Rightarrow P_f = 3730 \text{ Pa}$$

Other Solution :

As volume is constant

$$\therefore P \propto T$$

Since T is doubled (must be in Kelvin) so pressure must be doubled.


$$\therefore P_f = 2P_i$$

$$P_f = 2 \times 3.23 = 6.46 \text{ KPa}$$

$$P_f = 6460 \text{ Pa}$$

48. A voltage regulating circuit consisting of Zener diode, having break-down voltage of 10 V and maximum power dissipation of 0.4 W, is operated at 15 V. The approximate value of protective resistance in this circuit is _____ Ω .

Ans. (125)




$$P_D = 0.4W = 10i$$

$$i = 0.04A$$


$$R = \frac{15 - 10}{0.04} = \frac{5}{0.04} = 125 \Omega$$

- 49.** In the given figure the blocks A, B and C weigh 4 kg, 6 kg and 8 kg respectively. The co-efficient of sliding friction between any two surfaces is 0.5. The force \vec{F} required to slide the block C with constant speed is _____ N. (Used $g = 10 \text{ m/s}^2$)



Ans. (210)

Sol. For 8kg to move with constant velocity $F_{\text{net}} = 0$.



$$\therefore F = 90 + T + 50 \text{ (for 8kg block)}$$

$$T = 20 + 50 \text{ (for 6kg block)}$$


$$\therefore F = 210 \text{ N.}$$

- 50.** Sixty four rain drops of radius 1 mm each falling down with a terminal velocity of 10 cm/s coalesce to form a bigger drop. The terminal velocity of bigger drop is _____ cm/s.

Ans. (160)

Sol. $V_T = \frac{2r^2g}{9\eta}[\sigma - \rho]$

$$V_T \propto r^2$$



64 drop

$$64 \left(\frac{4}{3} \pi R_1^3 \right) = \frac{4}{3} \pi R_2^3$$

$$R_2 = 4R_1$$

$$\frac{(V_T)_1}{(V_T)_2} = \left(\frac{R_1}{R_2} \right)^2 = \left(\frac{1}{4} \right)^2$$

$$\frac{10}{(V_T)_2} = \frac{1}{16}$$

$$(V_T)_2 = 160 \text{ cm/sec}$$

CHEMISTRY

SECTION-A

51. Given below are two statements :

Statement-I Hybridisation, shape and spin only magnetic moment of $K_3[Co(CO_3)_3]$ is sp^3d^2 , octahedral and 4.9 BM respectively.

Statement-II Geometry, hybridisation and spin only magnetic moment values (BM) of the ions $[Ni(CN)_4]^{2-}$, $[MnBr_4]^{2-}$ and $[CoF_6]^{3-}$ respectively are square planar, tetrahedral, octahedral : dsp^2 , sp^3 , sp^3d^2 and 0, 5.9, 4.9.

In the light of the above statements, choose the **correct** answer from the options given below

- (1) Both statement-I and statement-II are false
- (2) Statement I is false but statement-II is true
- (3) Both statement-I and statement-II are true
- (4) Statement-I is true but statement-II is false

Ans. (3)

Sol. In $K_3[Co(CO_3)_3] \Rightarrow sp^3d^2$ hybridized, octahedral

\Rightarrow 4 unpaired electron

\Rightarrow 4.9 B.M.

$[Ni(CN)_4]^{2-} \Rightarrow dsp^2$ hybridized, square planar

\Rightarrow 0 unpaired electron

\Rightarrow 0 B.M.

$[MnBr_4]^{2-} \Rightarrow sp^3$ hybridized, tetrahedral

\Rightarrow 5 unpaired electron

\Rightarrow 5.9 B.M.

$[CoF_6]^{3-} \Rightarrow sp^3d^2$ hybridized, octahedral

\Rightarrow 4 unpaired electron

\Rightarrow 4.9 B.M.





52. $A \rightarrow D$ is an endothermic reaction occurring in three steps (elementary).

(i) $A \rightarrow B \Delta H_i = +ve$

(ii) $B \rightarrow C \Delta H_{ii} = -ve$


(iii) $C \rightarrow D \Delta H_{iii} = -ve$

Which of the following graphs between potential energy (y-axis) vs reaction coordinate (x-axis) correctly represents the reaction profile of $A \rightarrow D$?







Ans. (3)


Sol. Given :



Mechanism



53. Arrange the following alkenes in decreasing order of stability.




Choose the **correct** answer from the options given below :

- (1) III > I > II > IV (2) III > II > I > IV
(3) I > III > II > IV (4) I > III > IV > II

Ans. (3)

Sol. Stability order :



Trans is more stable than cis

54. Given below are statements about some molecules/ions.

Identify the **CORRECT** statements.

- A. The dipole moment value of NF_3 is higher than that of NH_3 .
B. The dipole moment value of BeH_2 is zero.
C. The bond order of O_2^{2-} and F_2 is same.
D. The formal charge on the central oxygen atom of ozone is -1 .
E. In NO_2 , all the three atoms satisfy the octet rule, hence it is very stable.

Choose the **correct** answer from the options given below :

- (1) A, B, C, D & E (2) B & C only
(3) B, C & D only (4) A, C & D only

Ans. (2)

- Sol. (A) Dipole moment : $\text{NF}_3 < \text{NH}_3$
(B) BeH_2 is 'sp' hybridized, linear molecule with zero dipole moment.
(C) $\text{O}_2^{-2} \Rightarrow$ bond order = 1
 $\text{F}_2 \Rightarrow$ bond order = 1
(D) Formal charge on central oxygen atom in O_3 is $+1$.
(E) In NO_2 ; nitrogen does not follow octet rule.

55. A solution is prepared by dissolving 0.3 g of a non-volatile non-electrolyte solute 'A' of molar mass 60 g mol^{-1} and 0.9 g of a non-volatile non-electrolyte solute 'B' of molar mass 180 g mol^{-1} in 100 mL H_2O at 27°C . Osmotic pressure of the solution will be

[Given : $R = 0.082 \text{ L atm K}^{-1} \text{ mol}^{-1}$]

- (1) 1.23 atm (2) 2.46 atm
(3) 0.82 atm (4) 1.47 atm

Ans. (2)

Sol. Mass of solute 'A' = 0.3 g

$$\text{Moles of solute 'A'} = \frac{0.3\text{g}}{60\text{g/mol}} = \frac{1}{200} \text{ mol}$$

Mass of solute 'B' = 0.9 g

$$\text{Moles of solute 'B'} = \frac{0.9 \text{ gm}}{180 \text{ g/mol}} = \frac{1}{200} \text{ mol}$$

Total molarity of all solutes

$$= \frac{2/200}{100} \times 1000 = \frac{1}{10} \text{ M}$$

$$\therefore \pi = \frac{1}{10} \times 0.082 \times 300$$

$$\pi = 2.46 \text{ atm.}$$


56. Among the following, the CORRECT combinations are :

- A. $\text{IF}_3 \rightarrow \text{T-shaped (sp}^3\text{d)}$
- B. $\text{IF}_5 \rightarrow \text{Square pyramidal (sp}^3\text{d}^2)$
- C. $\text{IF}_7 \rightarrow \text{Pentagonal bipyramidal (sp}^3\text{d}^3)$
- D. $\text{ClO}_4^- \rightarrow \text{Square planar (sp}^2\text{d)}$

Choose the **correct** answer from the options given below :


- (1) A, B and C only
- (2) A and B only
- (3) A, B, C and D
- (4) B, C and D Only

Ans. (1)




Hybridisation sp^3d (T-shape) Hybridisation sp^3d^2 (Square pyramidal)

Sol.



Hybridisation sp^3d^3 (Pentagonal bipyramidal) Hybridisation sp^3 (Tetrahedral)


57. The correct stability order of the following diazonium salts is



- (1) $\text{A} > \text{B} > \text{C} > \text{D}$
- (2) $\text{C} > \text{D} > \text{B} > \text{A}$
- (3) $\text{A} > \text{C} > \text{D} > \text{B}$
- (4) $\text{C} > \text{A} > \text{D} > \text{B}$

Ans. (3)

Sol. Correct order of stability



+M -M -M
EDG increases stability
EWG decreases stability

58. Consider a mixture 'X' which is made by dissolving 0.4 mol of $[\text{Co}(\text{NH}_3)_5\text{SO}_4] \text{Br}$ and 0.4 mol of $[\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$ in water to make 4 L of solution. When 2 L of mixture 'X' is allowed to react with excess of AgNO_3 , it forms precipitate 'Y'. The rest 2 L of mixture 'X' reacts with excess BaCl_2 to form precipitate 'Z'. Which of the following statements is **CORRECT**.

- (1) 0.2 mol of 'Z' is formed
- (2) 'Y' is BaSO_4 and 'Z' is AgBr
- (3) 0.4 mol of 'Z' is formed
- (4) 0.1 mol of 'Y' is formed

Ans. (1)

Sol. 0.4 mol $[\text{Co}(\text{NH}_3)_5\text{SO}_4]\text{Br} + 0.4 \text{ mol } [\text{Co}(\text{NH}_3)_5\text{Br}]\text{SO}_4$ are present in 4 lit. solution.

2 lit. of mixture will contain 0.2 mol of each complex.

2 lit. mixture on reaction with excess AgNO_3 , 0.2 mole AgBr will be formed (Y).

2 lit. mixture on reaction with excess BaCl_2 , 0.2 mole BaSO_4 will be formed (Z).

59. Given below are two statements

Statements-I : The number of paramagnetic species among $[\text{CoF}_6]^{3-}$, $[\text{TiF}_6]^{3-}$, V_2O_5 and $[\text{Fe}(\text{CN})_6]^{3-}$ is 3.


Statement-II : $\text{K}_4[\text{Fe}(\text{CN})_6] < \text{K}_3[\text{Fe}(\text{CN})_6] < [\text{Fe}(\text{H}_2\text{O})_6]\text{SO}_4 \cdot \text{H}_2\text{O} < [\text{Fe}(\text{H}_2\text{O})_6]\text{Cl}_3$ is the correct order in terms of number of unpaired electron(s) in the complexes.

In the light of the above statements, choose the **correct** answer from the options given below.

- (1) Both statement-I and statement-II are true
- (2) Both statement-I and statement-II are false
- (3) Statement-I is true but statement-II is false
- (4) Statement-I is false but statement-II is true

Ans. (1)

Sol. Paramagnetic species :



Diamagnetic species : V_2O_5

In $\text{K}_4[\text{Fe}(\text{CN})_6] \Rightarrow$ No. of unpaired electron = 0

$\text{K}_3[\text{Fe}(\text{CN})_6] \Rightarrow$ No. of unpaired electron = 1

$[\text{Fe}(\text{H}_2\text{O})_6]\text{SO}_4 \cdot \text{H}_2\text{O} \Rightarrow$ No. of unpaired electron = 4

$[\text{Fe}(\text{H}_2\text{O})_6]\text{Cl}_3 \Rightarrow$ No. of unpaired electron = 5

60. Consider three metal chlorides x, y and z, where x is water soluble at room temperature, y is sparingly soluble in water at room temperature and z is soluble in hot water. x, y and z are respectively

- (1) MgCl_2 , AgCl and AlCl_3
- (2) AgCl , Hg_2Cl_2 and PbCl_2
- (3) AlCl_3 , PbCl_2 and BaCl_2
- (4) CuCl_2 , AgCl and PbCl_2


Ans. (4)

Sol. MgCl_2 , AlCl_3 , CuCl_2 are water soluble at room temperature.

AgCl , Hg_2Cl_2 are sparingly soluble in water

PbCl_2 is soluble in hot water.

61. Match the List-I with List-II

List-I (Chloro derivative)		List-II (Example)	
A.	Vinyl Chloride	I.	$\text{CH}_2 = \text{CH} - \text{CH}_2\text{Cl}$
B.	Benzyl chloride	II.	$\text{CH}_3 = \text{CH}(\text{Cl})\text{CH}_3$
C.	Alkyl chloride	III.	$\text{CH}_2 = \text{CHCl}$
D.	Allyl chloride	IV.	

Choose the correct answer from the options given below :

- (1) A-IV, B-I, C-III, D-II
- (2) A-III, B-IV, C-I, D-II
- (3) A-III, B-IV, C-II, D-I
- (4) A-I, B-II, C-IV, D-III

Ans. (3)

Sol. Common Name (Theory based)

62. 'W' g of a non-volatile electrolyte solid solute of molar mass 'M' g mol^{-1} when dissolved in 100 mL water, decreases vapour pressure of water from 640 mm Hg to 600 mm Hg. If aqueous solution of the electrolyte boils at 375 K and K_b for water is $0.52 \text{ K kg mol}^{-1}$, then the mole fraction of the electrolyte solute (x_2) in the solution can be expressed as

(Given : density of water = 1 g/mL and boiling point of water = 373 K)

- (1) $\frac{1.3}{8} \times \frac{W}{M}$
- (2) $\frac{16}{2.6} \times \frac{W}{M}$
- (3) $\frac{2.6}{16} \times \frac{M}{W}$
- (4) $\frac{1.3}{8} \times \frac{M}{W}$

Ans. (1)

Sol. $P^\circ = 640 \text{ mm Hg}$

$P_s = 600 \text{ mm Hg}$

$\Delta P = 40 \text{ mm Hg}$

$$\text{moles of solute} = \frac{W}{M}$$

$$\frac{\Delta P}{P^0} = i \cdot X_{\text{solute}}$$

Again :

$$\Delta T_b = i \times k_b \times m$$

$$2 = i \times 0.52 \times \frac{W/M}{100} \times 1000$$

$$i = \frac{2}{5.2} \times \frac{M}{W}$$

$$X_{\text{solute}} = \frac{40}{640} \times \frac{1}{i} = \frac{1}{16} \times \frac{5.2}{2} \times \frac{W}{M}$$

$$X_{\text{solute}} = \frac{1.3}{8} \times \frac{W}{M}$$

63. Match the List-I with List-II

List-I (Isothermal process for ideal gas system)		List-II Work done ($V_f > V_i$)	
A.	Reversible expansion	I.	$w = 0$
B.	Free expansion	II.	$w = -nRT \ln \frac{V_f}{V_i}$
C.	Irreversible expansion	III.	$w = -P_{\text{ex}} (V_f - V_i)$
D.	Irreversible compression	IV.	$w = -P_{\text{ex}} (V_i - V_f)$

Choose the **correct** answer from the options given below :

- (1) A-IV, B-I, C-III, D-II
 (2) A-IV, B-II, C-III, D-I
 (3) A-I, B-III, C-II, D-IV
 (4) A-II, B-I, C-III, D-IV

Ans. (4)

Sol. (A) $W_{\text{Rev.}} = - \int P_{\text{gas}} dV$

$$W_{\text{Rev. Isot. Exp.}} = -nRT \ln \left[\frac{V_f}{V_i} \right]$$

(A) \rightarrow (II)

(B) Free expansion

$$W_{\text{irrev.}} = -P_{\text{ext}} \Delta V$$

$$P_{\text{ext}} = 0$$

$$W = 0$$

(B) \rightarrow (I)

(C) Irreversible expansion

$$W_{\text{irrev.}} = -P_{\text{ext}} \Delta V$$

$$W_{\text{irrev.}} = -P_{\text{ext}} (V_f - V_i)$$


(C) \rightarrow (III)

(D) Irreversible compression

$$W_{\text{irrev.}} = -P_{\text{ext}} \Delta V$$

$$W_{\text{irrev.}} = -P_{\text{ext}} (V_i - V_f)$$


64. A student is given one compound among the following compounds that gives positive test with Tollen's reagent.



The compound is :

- (1) D (2) A
 (3) B (4) C


Ans. (4)



65. A hydroxy compound (X) with molar mass 122 g mol^{-1} is acetylated with acetic anhydride, using a large excess of the reagent ensuring complete acetylation of all hydroxyl groups. The product obtained has a molar mass of 290 g mol^{-1} . The number of hydroxyl groups present in compound (X) is :

- (1) 3 (2) 5
 (3) 2 (4) 4

Ans. (4)



$$\text{No. of OH groups} = \frac{290 - 122}{42} = 4$$

66. At 27°C in presence of a catalyst, activation energy of a reaction is lowered by 10 kJ mol⁻¹. The

logarithm ratio of $\frac{k(\text{catalysed})}{k(\text{uncatalysed})}$ is

(Consider that the frequency factor for both the reactions is same)

- (1) 17.41
- (2) 1.741
- (3) 3.482
- (4) 0.1741

Ans. (2)

Sol.
$$\frac{K_{\text{catalyst}}}{K_{\text{uncatalyst}}} = e^{\frac{\Delta E_a}{RT}}$$

$$\ln \frac{K_{\text{catalyst}}}{K_{\text{uncatalyst}}} = \frac{\Delta E_a}{RT}$$

$$\log \frac{K_{\text{catalyst}}}{K_{\text{uncatalyst}}} = \frac{\Delta E_a}{2.303RT}$$


$$= \frac{10 \times 1000}{2.303 \times 8.314 \times 300}$$

$$\log \frac{K_{\text{catalyst}}}{K_{\text{uncatalyst}}} = 1.741$$

67. Given below are two statements :

Statement I : 'C-Cl' bond is stronger in CH₂ = CH - Cl than CH₃ - CH₂ - Cl

Statement II : The given optically active




solution that can rotate the plane polarized light.

In the light of the above statements, choose the **correct** answer from the options given below

- (1) Statement I is false but Statement II is true
- (2) Both Statement I and Statement II are true
- (3) Both Statement I and Statement II are false
- (4) Statement I is true but Statement II is false


Ans. (4)

Sol. **Statement-I :**





C-Cl bond is strong in vinyl chloride because of double bond character.

Statement-II :



Racemic mixture is optically inactive, which can not rotate PPL.

68. Consider the following two reactions A and B.



Numerical value of [molar mass of x + molar mass of y] is _____.






- (1) 4
- (2) 88
- (3) 46
- (4) 160

Ans. (3)

Sol. x = H₂ (gas), y = CO₂ (gas)

Sum of molar mass = 2 + 44 = 46

69. Arrange the following carbanions in the decreasing order of stability



Choose the **correct** answer from the options given below :

- (1) I > II > IV > V > III
- (2) I > IV > II > V > III
- (3) IV > I > II > V > III
- (4) IV > II > I > III > V

Ans. (3)

Sol. Electron withdrawing groups increases stability of carbanion and electron donating groups decreases stability of carbanion.

70. Given below are two statements :

Statement I : $\text{K} > \text{Mg} > \text{Al} > \text{B}$ is the correct order in terms of metallic character.

Statement II : Atomic radius is always greater than the ionic radius for any element.

In the light of the above statements, choose the **correct** answer from the options given below

- (1) Both Statement I and Statement II are true
- (2) Both Statement I and Statement II are false
- (3) Statement I is false but Statement II is true
- (4) Statement I is true but Statement II is false

Ans. (4)



Sol. Metallic character of s-block elements is greater than p-block elements.

Anionic radius is greater than atomic radius but cationic radius is always less than atomic radius for any element.

SECTION-B

71. Electricity is passed through an acidic solution of Cu^{2+} till all the Cu^{2+} was exhausted, leading to the deposition of 300 mg of Cu metal. However, a current of 600 mA was continued to pass through the same solution for another 28 minutes by keeping the total volume of the solution fixed at 200 mL. The total volume of oxygen evolved at STP during the entire process is _____ mL. (Nearest integer)

[Given :



Molar mass of Cu = 63.54 g mol⁻¹

Molar mass of O₂ = 32 g mol⁻¹

Faraday Constant = 96500 C mol⁻¹

Molar volume at STP = 22.4 L]

Ans. (111)

Sol. Eq of Cu = Eq of O₂

$$\frac{300 \times 10^{-3} \times 2}{63.54} = n_{\text{O}_2} \times 4$$

$$2.36 \times 10^{-3} = n_{\text{O}_2}$$

When current is further passed

$$n_{\text{O}_2} \times 4 = \frac{600 \times 28 \times 60}{96500 \times 1000}$$

$$n_{\text{O}_2} = 2.611 \times 10^{-3}$$

Total O₂ released


$$= [10^{-3} \times (2.36 + 2.611)] \times 22400 \text{ ml}$$

$$= 111.35 \text{ ml}$$


72. Consider two Group IV metal ions X²⁺ and Y²⁺.

A solution containing 0.01 MX²⁺ and 0.01 MY²⁺ is saturated with H₂S. The pH at which the metal sulphide YS will form as a precipitate is _____ (Nearest integer)

(Given :




Ans. (4)



For precipitation of $XS(s)$

$$[X^{+2}][S^{2-}] \geq K_{sp}(XS)$$


$$[S^{2-}] \geq \frac{1 \times 10^{-22}}{0.01} = 10^{-20}$$



For precipitation of $YS(s)$

$$[Y^{+2}][S^{2-}] \geq K_{sp}(YS)$$

$$[S^{2-}] \geq \frac{4 \times 10^{-16}}{10^{-2}} = 4 \times 10^{-14}$$



$$\frac{[S^{2-}][H^+]^2}{H_2S} = K_{a1} \times K_{a2} = 1 \times 10^{-21}$$

$$[S^{2-}] = \frac{1 \times 10^{-21} \times [H_2S]}{[H^+]^2} \geq 4 \times 10^{-14}$$

$$[H^+]^2 \leq \frac{1}{4} \times 10^{-7} \times 10^{-1}$$

$$[H^+] \leq \frac{1}{2} \times 10^{-4} \Rightarrow \text{pH} \geq 4.3$$

73. The hydrogen spectrum consists of several spectral lines in Lyman series (L_1, L_2, L_3, \dots ; L_1 has lowest energy among Lyman series). Similarly it consists of several spectral lines in Balmer series (B_1, B_2, B_3, \dots ; B_1 has lowest energy among Balmer lines). The energy of L_1 is x times the energy of B_1 . The value of x is _____ $\times 10^{-1}$ (Nearest integer)

Ans. (54)

Sol. $\Delta E(L_1) = 13.6 \times Z^2 \left(\frac{1}{1^2} - \frac{1}{2^2} \right) = 13.6Z^2 \times \frac{3}{4}$

$$\Delta E(B_1) = 13.6 \times Z^2 \times \left(\frac{1}{2^2} - \frac{1}{3^2} \right) = 13.6 \times Z^2 \times \frac{5}{4 \times 9}$$

$$\frac{\Delta E(L_1)}{\Delta E(B_1)} = \frac{3}{5} \times 9 = \frac{27}{5} = x$$

$$x = \left(\frac{27}{5} \times 10 \right) \times 10^{-1} = 54 \times 10^{-1}$$

74. In Dumas method for estimation of nitrogen, 0.50 g of an organic compound gave 70 mL of nitrogen collected at 300 K and 715 mm pressure. The percentage of nitrogen in the organic compound is _____ %

(Aqueous tension at 300 K is 15 mm).

Ans. (15)

Sol. $P_{N_2} = (715 - 15) \text{ mm} = \frac{700}{760} \text{ atm}$

$$V_{N_2} = 70 \text{ ml} = \frac{70}{1000} \text{ l}$$

$$n_{N_2} = \frac{PV}{RT} = \frac{\left(\frac{700}{760} \right) \times \left(\frac{70}{1000} \right)}{0.0821 \times 300}$$


$$W_{N_2} = \frac{700}{760} \times \frac{70}{0.0821 \times 300} \times 28$$

$$\% N = \frac{W_{N_2}}{0.5} \times 100 = \frac{700}{760} \times \frac{70/1000}{0.0821 \times 300} \times 28 \times 100$$


$$= 14.65\% \approx 15$$

75. X and Y are the number of electrons involved, respectively during the oxidation of I^- to I_2 and S^{2-} to S by acidified $K_2Cr_2O_7$. The value of $X + Y$ is _____.

Ans. (12)



no. of moles e^- involved = $x = 6$



No. of moles e^- involved = $y = 6$

Sum of $x + y = 6 + 6 = 12$